

AL.2.1998-14

C.2.

CANADIANA

DEC 22 1997

ECOLOGICAL LAND CLASSIFICATION  
AND  
VEGETATION DESCRIPTION  
OF  
UPPER BOB CREEK  
ECOLOGICAL RESERVE



**Alberta**  
ENVIRONMENTAL  
PROTECTION





## ACKNOWLEDGEMENTS

The following people participated in 1994 and 1995 field work in the Upper Bob Creek Ecological Reserve. From Resource Data Division - Dave Downing, Ed Kopyck, AJ Brown, and Tracy Rynick. From Alberta Forest Service - Tracy Rynick and Angela Ragan.

I would like to thank Mike Willoughby of the Alberta Forest Service for his friendly discussions and helpful advice. Vegetation cover data from the Upper Bob Creek field plots were included in the following vegetation community type analysis by Willoughby et al (1996) and their classification is used in this report.

The base map and final map were compiled by AJ Brown of Resource Data Division.

The project was coordinated by Beth Cornish. In addition I would like to acknowledge the assistance of Ed Kopyck, Ed Brown, and Mike Alexander, Alberta Forest Service, Sherwood.

## ECOLOGICAL LAND CLASSIFICATION

### AND

## VEGETATION DESCRIPTION

### OF

## UPPER BOB CREEK

## ECOLOGICAL RESERVE

Beth Cornish  
Resource Data Division  
Alberta Environmental Protection  
Edmonton, Alberta

February 1997



## ACKNOWLEDGEMENTS

The following people participated in 1994 and 1995 field work in the Upper Bob Creek Ecological Reserve: from Resource Data Division - Dave Downing, Ed Karpuk, Al Benner, and Troy Berreth; from Alberta Forest Service - Tracy Rainforth and Angela Bogan.

I would like to thank Mike Willoughby of the Alberta Forest Service for his friendly discussions and helpful advice. Vegetation cover data from the Upper Bob Creek field plots were included in the Montane vegetation community type analysis by Willoughby et al (1996), and their classification is used in this report.

The base map, and final drafts of the ecological land classification and special features maps, were completed by Al Benner, with GIS support from Lucy Faferek, both of Resource Data Division.

The project was coordinated by Ed Karpuk, Resource Data Division. In addition I would like to acknowledge the assistance of Russ Wells, Resource Data Division, Edmonton, and Mike Alexander, Alberta Forest Service, Blairmore.



Digitized by the Internet Archive  
in 2017 with funding from  
University of Alberta Libraries

<https://archive.org/details/ecologicalandcl1997corn>

# TABLE OF CONTENTS

|                                                              |    |
|--------------------------------------------------------------|----|
| ACKNOWLEDGEMENTS .....                                       | i  |
| 1.0 INTRODUCTION .....                                       | 1  |
| 2.0 STUDY AREA DESCRIPTION .....                             | 1  |
| 2.1 Location .....                                           | 1  |
| 2.2 Climate .....                                            | 3  |
| 2.3 Hydrography .....                                        | 3  |
| 2.4 Physiography and Bedrock Geology .....                   | 3  |
| 2.5 Surficial Geology and Parent Materials .....             | 4  |
| 2.6 Soils .....                                              | 4  |
| 2.7 Vegetation .....                                         | 4  |
| 3.0 METHODS .....                                            | 5  |
| 3.1 Field Survey .....                                       | 5  |
| 3.2 Data Analysis .....                                      | 5  |
| 3.3 Ecological Land Classification and Mapping .....         | 5  |
| 3.4 Polygon Database and Legend .....                        | 6  |
| 3.5 Special Features Map .....                               | 7  |
| 4.0 RESULTS .....                                            | 7  |
| 4.1 Ecological Land Classification .....                     | 7  |
| 4.2 Vegetation Community Types .....                         | 7  |
| 4.2.1 Rough fescue-Idaho fescue-Parry oatgrass .....         | 8  |
| 4.2.2 Idaho fescue - Parry oatgrass - Rough fescue .....     | 8  |
| 4.2.3 Rough fescue - Kentucky bluegrass - Timothy .....      | 9  |
| 4.2.4 Kentucky bluegrass - Rough fescue .....                | 10 |
| 4.2.5 Pine grass - Hairy wild rye / Wild strawberry .....    | 11 |
| 4.2.6 Bog birch - Basket willow - Myrtle-leaved willow ..... | 12 |
| 4.2.7 Aspen / Bearberry / Rough fescue .....                 | 14 |
| 4.2.8 Aspen / Rose / Pine grass .....                        | 15 |
| 4.2.9 Aspen - White spruce/ Blueberry .....                  | 16 |
| 4.2.10 Spruce - Balsam poplar / Snowberry .....              | 17 |
| 4.2.11 Douglas fir /Hairy wild rye .....                     | 18 |
| 4.2.12 Lodgepole pine/ White meadowsweet .....               | 19 |
| 4.2.13 White spruce- Douglas fir/ White meadowsweet .....    | 20 |
| 4.2.14 White spruce/moss .....                               | 21 |
| 4.2.15 White spruce/ Thimbleberry .....                      | 22 |
| 4.2.16 Limber pine - Douglas fir/Juniper - Bearberry .....   | 23 |
| 4.3 Significant Plant Species .....                          | 23 |



|                                                                                                                                                      |    |
|------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 5.0 LITERATURE CITED .....                                                                                                                           | 27 |
| APPENDIX A. Map polygon database.....                                                                                                                | 29 |
| APPENDIX B. Descriptions of physical land unit codes from Appendix A .....                                                                           | 35 |
| APPENDIX C. Summary of vegetation, landform and soils information collected at each site,<br>Upper Bob Creek Ecological Reserve, 1994 and 1995. .... | 37 |
| APPENDIX D. Significant plant species which may occur in the Ecological Reserve, and status,<br>habitat and occurrence details. ....                 | 40 |
| APPENDIX E. Ranking system used to evaluate status of significant plant species. ....                                                                | 42 |

## LIST OF TABLES

|                                                                                       |    |
|---------------------------------------------------------------------------------------|----|
| Table 1. Rare or uncommon plants recorded at Upper Bob Creek Ecological Reserve. .... | 25 |
|---------------------------------------------------------------------------------------|----|

## LIST OF FIGURES AND MAPS

|                                                                  |   |
|------------------------------------------------------------------|---|
| Fig. 1. Location of the Upper Bob Creek Ecological Reserve. .... | 2 |
|------------------------------------------------------------------|---|

MAP 1. Ecological land classification of Upper Bob Creek Ecological Reserve.

MAP 2. Significant vegetation features at Upper Bob Creek Ecological Reserve.



## **1.0 INTRODUCTION**

The Upper Bob Creek Ecological Reserve (Fig. 1) was formally designated in 1987 under the Wilderness Areas, Ecological Reserves and Natural Areas Act. It includes representative montane landscapes which support a diverse biota, including a number of rare and uncommon plants and animals. The area includes important elk wintering grounds (O'Leary et al 1989), has a long history of cattle grazing and is a popular recreation area.

To facilitate resource management in the Ecological Reserve, Alberta Land and Forest Service requested a study to gather baseline information on vegetation resources and their relation to landscape characteristics. This information would be used as an aid in evaluating the condition of the natural resource, and for wildlife habitat management and domestic grazing management.

Objectives of the study were to:

1. Prepare an ecological land classification map, showing vegetation types matched to physical land classification units.
2. Prepare a description of the plant communities that occur in the study area based on the 1994 and 1995 field work, with reference to previously defined vegetation types or ecosite phases (Willoughby et al 1996, and Archibald et al 1996).
3. Prepare a map and list of significant vegetation features.

## **2.0 STUDY AREA DESCRIPTION**

### **2.1 Location**

The study area was limited to the Upper Bob Creek Ecological Reserve, located in southwest Alberta, about 70 km north of the town of Pincher Creek (Fig. 1). It is situated in the foothills east of the Livingstone Range, north of the Oldman River. Most of the eastern and all of the western boundaries of the Reserve are defined by the watershed of the upper reaches of Bob Creek. The watershed divide on the top of the north-south trending Whaleback Ridge forms the eastern boundary, and the western boundary similarly follows the watershed divide. The south end of Township 12 forms the southern boundary of the Reserve. The northern boundary is irregular, following legal survey gridlines, the watershed divide between Chaffen Creek and Bob Creek, and the watercourse of an unnamed creek.

The Upper Bob Creek Ecological Reserve covers approximately 2600 ha (6427 acres) and includes the following land areas:

- in Twp 12 Rge 2 W5M: Sections 4-8, and 17; and parts of Sections 9, 16, 18-21, 28, 29;
- in Twp 12 Rge 3 W5M: small portions of Sections 1, 12 and 13, east of the height of land.






Fig. 1. Location of the Upper Bob Creek Ecological Reserve.



## **2.2 Climate**

The Upper Bob Creek Ecological Reserve occurs in the Montane Natural Subregion within the Rocky Mountain Natural Region (Achuff 1994). Weather data collected in an area south of, and perhaps warmer than, the study area, indicated an average annual precipitation of about 600 mm (Coleman and Beaver Mines data - Environment Canada 1982; Achuff 1994). The area is characterized by fairly warm summers (mean temperature May to September of about 12° C) and cool winters (mean January temperature of -8° C) moderated by frequent chinooks (Achuff 1994).

Wind represents a significant climatic factor influencing vegetation. Fairly strong southwesterly to westerly winds occur more than 40% of the time (Strong 1979; Brown et al 1986). The west winds are especially prevalent from October to February. Chinook winds (with temperatures above 4 °C) occur an average of 25 or more days per year in the study area (Strong 1979). Strong winds affect vegetation primarily through physical damage, increased evapotranspiration and redistribution of snow to the lee sides of the hill crests.

## **2.3 Hydrography**

Most of the Ecological Reserve is drained by the upper 5 km of Bob Creek, a perennial stream which flows to the south and eventually into the Oldman River. A small section of the north end of the study area drains into Chaffin Creek. The headwaters of Bob Creek are represented by several small tributaries which flow off the dissected ridges on the north, east and west sides of the Reserve. Flow in some of the tributary creeks may be intermittent during dry periods (Leskiw 1993). Seepage areas are scattered throughout the catchment basins of these tributaries.

## **2.4 Physiography and Bedrock Geology**

The terrain includes many dissected ridges and a wide creek valley. Elevation ranges from about 1460 m in the valley bottom at the south end of the study area, to 1909 m at the highest ridgetops along the western boundary. The highest elevation along the eastern boundary is about 1750 m. Slopes vary from nearly flat, such as in poorly drained areas on the valley floor, to 70% or more on the upper slopes.

The Ecological Reserve occurs within the Southern Foothills District of the Rocky Mountain Foothills Region (Pettapiece 1986). A band of north-south trending ridges occurs in the study area, representing a transition zone between the Rocky Mountains and the rolling topography of the Porcupine Hills to the east. Bedrock formations represented in the study area are of Upper Cretaceous age. They include thick-bedded sandstones of the Brazeau Formation on the ridges, and a formation series called the Alberta Group in the valleys (Green 1972). Physiographic features of the Ecological Reserve are summarized in greater detail in Leskiw (1993).



## 2.5 Surficial Geology and Parent Materials

Colluvial veneers and residuum (weathered bedrock) occur on the ridges and upper slopes, overlying the Brazeau Formation bedrock. On the lower slopes, morainal blankets and veneers are dominant, along with lesser amounts of colluvial veneer over the inclined bedrock of the Alberta Group formations. In the bottomlands of the valleys of Bob Creek and its larger tributaries, undulating morainal blankets occur with a veneer of recent fluvial deposits (Leskiw 1993).

## 2.6 Soils

Rapidly drained shallow Lithic and Orthic Eutric Brunisols have developed on the weathered bedrock on the ridges, with some rapid to very rapidly drained Orthic Regosols on the most exposed ridgetops. On the upper slopes, well- to rapidly drained Orthic Eutric Brunisols predominate, with Orthic Gray Luvisols occurring less frequently. Orthic Black and, less commonly, Dark Gray Chernozems are found on the lower slopes. These medium-textured soils tend to be moderately well to well-drained. Black Chernozems have also formed on the fluvial and morainal deposits on the valley bottoms. Pockets of poorly drained to very poorly drained areas occur in seepage areas and along streams where Gleysols, most commonly Orthic Humic Gleysols, have developed.

## 2.7 Vegetation

The Montane Natural Subregion in Alberta is characterized by a mosaic of forest and grassland communities, and is distinguished from other subregions by the presence of Douglas fir (*Pseudotsuga menziesii*), limber pine (*Pinus flexilis*) and lodgepole pine (*Pinus contorta*) (Achuff 1994). Complex topography, varying from gently undulating valley floor to ridged and dissected upland, has created variable ecological conditions, controlled by factors such as elevation, aspect, soils and parent materials. Therefore, many different vegetation community types can be found even across small areas. There is a large amount of overlap between the forest and grassland in the Reserve and some community types are recognized as being transitional from one to the other.

Aspect is one of the most important site factors controlling vegetation in the study area. However, differences in vegetation related to aspect are more obvious in some areas of the study area than in others. At lower elevations east and west of Bob Creek and on the lower slopes of Whaleback Ridge, the effects of aspect are very pronounced. South-facing slopes are characterized by grasslands, with coniferous forest (dominated most commonly by lodgepole pine or a mix of lodgepole pine and Douglas fir) occurring on the cooler, moister north-facing slopes. The west-central part of the Ecological Reserve is, in contrast, heavily forested, despite variable aspects. However, as elevation increases on the ridge at the western boundary, larger areas of grassland are again found.



### **3.0 METHODS**

#### **3.1 Field Survey**

Site selection for the 1994 and 1995 range surveys was based on a pre-stratification of the area using landform and soil characteristics described by Leskiw (1993). Field data collection was conducted between June 14 and July 7 in 1994, and between August 1 and 13 in 1995. Twenty-three sites on primary and secondary range were surveyed in 1994 by crews from Resource Data Division and Alberta Forest Service (Fig. 1). In 1995, 18 forested sites were examined by a crew from Alberta Forest Service.

Vegetation surveys followed the procedure outlined in the Range Survey Manual (Alberta Forest Service 1990). At each site, a series of 15 microplots, 20 cm X 50 cm in size, was established along a 30 m transect in a representative part of the site. Within each microplot, all plant species were recorded and percent canopy cover estimated. Information was recorded on MF5 and MF6 Vegetation Inventory forms according to the guidelines in the Range Survey Manual. Physical site conditions were assessed at each site and recorded on LISD site forms, based on the methods in the Ecological Land Survey Site Description Manual (Alberta Environmental Protection 1994). At most sites, a soil pit was excavated in order to check parent material and classify local soil conditions.

#### **3.2 Data Analysis**

Vegetation data collected in 1994 and 1995 were analyzed using cluster analysis (SAS) and ordination (DECORANA, Gauch 1982) (Willoughby et al 1996). Sites were ranked based on their similarity to each other in terms of species cover values. Thus, sites of similar species composition were consolidated into derived groups or community types. Species composition of each newly identified community type was then described based on an average of the species cover values of the grouped sites.

Many of the vegetation community types found at the 41 sites examined in the study area corresponded to types described for the Montane Natural Subregion by other workers. These descriptions (Willoughby et al 1996) were based in part on the data collected in 1994 and 1995 in the Upper Bob Creek Ecological Reserve. Archibald et al (1996) outline Montane plant community types within a hierarchical ecological framework.

Common and scientific names for plant species are based on the Alberta Plants and Fungi Master Species List and Species Group Checklists (Alberta Environmental Protection 1993).

#### **3.3 Ecological Land Classification and Mapping**

A 1:15 000 scale ecological land classification map was created, matching existing physical land classification units to vegetation. This approach used a combination of landscape characteristics



(especially landform or parent material, soil type and slope) and vegetation composition to define map units. Thus, vegetation type information was overlaid on mapped landscape information.

The sources of information about landforms, slopes, parent materials and soil types in the Reserve were the physical land classification (PLC) completed on 1:20 000 scale aerial photographs (Leskiw 1993), and the site data collected in 1994 and 1995 (this study). Vegetation information was gained from two sources: the 1994 and 1995 field data, and, for forested polygons not visited in 1994 or 1995, from Alberta Vegetation Inventory (AVI). The AVI was based on interpretation of 1:20 000 scale aerial photography. This information was then synthesized to create mapped polygons representing ecological units.

Because of the variability in vegetation even across short distances in the study area, and the limitation of aerial photograph interpretation, it was often difficult to accurately predict a single vegetation community type in mapped polygons. For example, for areas not visited in the field, it was not possible to interpret forest understory forb species, nor to differentiate grazed from ungrazed grassland. These areas could not be described to vegetation community type level with sufficient accuracy based only on AVI information or aerial photograph interpretation. Therefore, predicted vegetation in each polygon was described to the somewhat more general Ecosite Phase level (Archibald et al 1996) for forested sites, and to a physiognomic type category (e.g. fescue grassland; wetland shrub) for non-forested sites.

Dominant overstory tree species define Ecosite Phases, which, with local environmental conditions, control species composition of the understory (Archibald et al 1996). Thus, vegetation described to the Ecosite Phase level can more easily be correlated to both AVI information and to site environmental conditions. An example of an Ecosite Phase found in the study area is "thimbleberry/pine grass Aw". This describes a site where the dominant overstory tree species is aspen (Aw). The characteristic understory species, thimbleberry and pine grass, are an expression of the site environmental conditions, namely a mesic moisture regime and a rich nutrient regime. Details of specific Ecosite Phases, including plant species composition, are provided in Archibald et al (1996).

### **3.4 Polygon Database and Legend**

A tabular polygon database was constructed, which serves as a map polygon legend. It describes the dominant vegetation, as well as certain landscape and soils characteristics for each polygon. Thus map polygon numbers were related to the following list of parameters, each of which was included as a separate field in the database:

Landform and surficial materials

Soil types = Soil1, Soil2

Moisture regime

Nutrient regime

Vegetation description - Ecosite Phase or physiognomic type



### **3.5 Special Features Map**

A second map, illustrating the location of populations of rare or uncommon plant species, was also produced at a scale of 1:15 000.

## **4.0 RESULTS**

### **4.1 Ecological Land Classification**

Map 1 shows the delineation of ecological units in the study area. Appendix A represents the polygon database, describing the vegetation and physical land characteristics (Appendix B) of each numbered polygon. The map legend includes vegetation type codes, based on ecosite phase or physiognomic type, and land unit codes, based on the predominant parent material in the polygon.

### **4.2 Vegetation Community Types**

Field data from the 41 sites were classified into 15 different vegetation community types, which can be grouped into the following major categories: five grassland types, one shrubland type, two deciduous forest types, two mixed forest types, and five coniferous forest types. Other vegetation community types which may occur in the Ecological Reserve but were not sampled in this study are described in Willoughby et al (1996) and Archibald et al (1996).

The plant community types sampled in the Ecological Reserve are described below. Scientific names of plant species discussed in the text are found in the accompanying species composition tables. A summary of vegetation community type, ecosite phase (for forested sites), parent materials and soils information from each site is given in Appendix C.

#### **Grassland community types**

Grassland communities are found on recent fluvial deposits in the Bob Creek valley, till or colluvial deposits on the south-facing lower and middle slopes of ridges, and thin rocky soils of the upper slopes. Five grassland community types were differentiated in the study area and their composition is governed by environmental conditions and grazing pressure. Transitional communities are also common, where types grade into each other or towards forest communities, as environmental conditions change.

These community types were found at specific locations in the Reserve. However, because the different grassland types could not be separated on aerial photographs, the distribution of the individual grassland communities could not be outlined in detail on Map 1. Field mapping of these communities is necessary.



#### 4.2.1 Rough fescue-Idaho fescue-Parry oatgrass

This community type was found at 13 sites examined in this study. It is the modal, or normal climax, grassland community type on undisturbed Black Chernozemic soils in the southern Alberta foothills (Willoughby et al 1996). In this study, it occurred primarily on Black Chernozems on the till and fluvial deposits at lower elevations. However, less commonly it was found on shallow Eutric Brunisols, on south-facing slopes at higher elevations. Rough fescue (*Festuca scabrella*), Parry oat grass (*Danthonia parryi*) and Idaho fescue (*Festuca idahoensis*) are the dominant grass species on modal sites not heavily grazed. There is a highly diverse forb layer, and important forb species include old man's whiskers (*Geum triflorum*), cut-leaved anemone (*Anemone multifida*) and sticky purple geranium (*Geranium viscosissimum*). With light grazing in this community, cover of Parry oat grass will increase (Johnston 1961; Moss and Campbell 1947). Rough fescue grasslands occur at elevations as high as 2121 m in southwest Alberta (Willoughby et al 1996), but in the Porcupine Hills, east of the study area, Jaques (1979) found that fescue grasslands occurred most extensively at elevations of 1500 m or less.

| Upper Bob Creek Community Type: Rough fescue - Idaho fescue - Parry oatgrass |                               |                                        |                  |
|------------------------------------------------------------------------------|-------------------------------|----------------------------------------|------------------|
| Sites: 1,3,5,6,8,10,12,13,20,23,24,26,28                                     |                               | Range classification: primary          |                  |
| Moisture regime: submesic to mesic                                           |                               | Soil drainage: well to moderately well |                  |
| Nutrient regime: mesotrophic<br>(submesotrophic to permesotrophic)           |                               | Soil subgroup: O.BL; O.EB              |                  |
| Slope (range): 0-60%                                                         |                               | Aspect: south                          |                  |
| Key and indicator species:                                                   |                               |                                        |                  |
| Scientific name                                                              |                               | Common name                            | Canopy cover (%) |
| Graminoids:                                                                  | <i>Festuca scabrella</i>      | Rough fescue                           | 29               |
|                                                                              | <i>Festuca idahoensis</i>     | Idaho fescue                           | 11               |
|                                                                              | <i>Danthonia parryi</i>       | Parry oat grass                        | 11               |
|                                                                              | <i>Carex obtusata</i>         | Blunt sedge                            | 8                |
|                                                                              |                               |                                        |                  |
| Forbs:                                                                       | <i>Geum triflorum</i>         | Old man's whiskers                     | 4                |
|                                                                              | <i>Penstemon confertus</i>    | Yellow beardtongue                     | 4                |
|                                                                              | <i>Anemone multifida</i>      | Cut-leaved anemone                     | 2                |
|                                                                              | <i>Geranium viscosissimum</i> | Sticky purple geranium                 | 2                |
|                                                                              | <i>Dodecatheon conjugens</i>  | Mountain shooting star                 | 1                |
| Shrubs:                                                                      | <i>Potentilla fruticosa</i>   | Shrubby cinquefoil                     | 3                |



#### 4.2.2 Idaho fescue - Parry oatgrass - Rough fescue

This community type was found at two lower slope sites in the study area. Compared to type #1 above, cover values of Idaho fescue and Parry oatgrass have increased in proportion to a decrease in rough fescue cover. Development of this community can be attributed to drier soil conditions or to grazing pressure, or a combination of both. Rough fescue is a decreaser (Wroe et al 1988), and cover values are reduced with grazing. In contrast, light grazing favours Idaho fescue and Parry oatgrass (Johnston 1961), as well as dryland sedge species such as blunt sedge (Moss and Campbell 1947). Soil conditions on grazed sites are often slightly warmer and drier than similar ungrazed sites, according to a study in the Porcupine Hills (Johnston 1961).

| Upper Bob Creek Community Type: Idaho fescue - Parry oatgrass - Rough fescue |                               |                                |                  |
|------------------------------------------------------------------------------|-------------------------------|--------------------------------|------------------|
| Sites: 2, 27                                                                 |                               | Range classification: Primary  |                  |
| Moisture regime: subxeric - submesic                                         |                               | Soil drainage: rapidly to well |                  |
| Nutrient regime: mesotrophic                                                 |                               | Soil subgroup: O.EB; O.BL      |                  |
| Slope % (range): 21 (0-36)                                                   |                               | Aspect: south                  |                  |
| Key and indicator species:                                                   |                               |                                |                  |
| Scientific name                                                              |                               | Common name                    | Canopy cover (%) |
| Graminoids:                                                                  | <i>Festuca idahoensis</i>     | Idaho fescue                   | 18               |
|                                                                              | <i>Festuca scabrella</i>      | Rough fescue                   | 10               |
|                                                                              | <i>Danthonia parryi</i>       | Parry oat grass                | 16               |
|                                                                              | <i>Carex obtusata</i>         | Blunt sedge                    | 3                |
|                                                                              | <i>Koeleria macrantha</i>     | June grass                     | 4                |
|                                                                              |                               |                                |                  |
| Forbs:                                                                       | <i>Achillea millefolium</i>   | Yarrow                         | 2                |
|                                                                              | <i>Geum triflorum</i>         | Old man's whiskers             | 3                |
|                                                                              | <i>Lupinus sericeus</i>       | Silky perennial lupine         | 4                |
|                                                                              | <i>Lithospermum ruderales</i> | Woolly gromwell                | 1                |
| Shrubs:                                                                      | <i>Potentilla fruticosa</i>   | Shrubby cinquefoil             | 2                |
|                                                                              | <i>Rosa acicularis</i>        | Prickly rose                   | 1                |

#### 4.2.3 Rough fescue - Kentucky bluegrass - Timothy

This site represents a rough fescue grassland which has been invaded by non-native species. It occurred on a lower slope just east of Bob Creek, in an area well utilized by cattle. The non-native species Kentucky bluegrass does well once it becomes established. The rhizomatous growth habit often allows it to eventually dominate the site. Timothy has been planted as a hay



crop outside the study area, and reproduces prolifically by seed.

| Upper Bob Creek Community Type: Rough fescue - Kentucky bluegrass - Timothy |                              |                               |                  |
|-----------------------------------------------------------------------------|------------------------------|-------------------------------|------------------|
| Sites: #29                                                                  |                              | Range classification: Primary |                  |
| Moisture regime: mesic                                                      |                              | Soil drainage: well           |                  |
| Nutrient regime: mesotrophic                                                |                              | Soil subgroup: O.BL           |                  |
| Slope % (range): 5%                                                         |                              | Aspect: west                  |                  |
| Key and indicator species:                                                  |                              |                               |                  |
| Scientific name                                                             |                              | Common name                   | Canopy cover (%) |
| Graminoids:                                                                 | <i>Festuca scabrella</i>     | Rough fescue                  | 16               |
|                                                                             | <i>Poa pratensis</i>         | Kentucky bluegrass            | 12               |
|                                                                             | <i>Phleum pratense</i>       | Timothy                       | 10               |
|                                                                             | <i>Festuca idahoensis</i>    | Idaho fescue                  | 15               |
|                                                                             | <i>Danthonia parryi</i>      | Parry oat grass               | 5                |
|                                                                             |                              |                               |                  |
| Forbs:                                                                      | <i>Achillea millefolium</i>  | Yarrow                        | 4                |
|                                                                             | <i>Geum triflorum</i>        | Old man's whiskers            | 20               |
|                                                                             | <i>Taraxacum officinale</i>  | Dandelion                     | 6                |
|                                                                             | <i>Dodecatheon conjugens</i> | Shooting star                 | 12               |
| Shrubs:                                                                     | <i>Potentilla fruticosa</i>  | Shrubby cinquefoil            | 4                |

#### 4.2.4 Kentucky bluegrass - Rough fescue

This community type also represents a rough fescue grassland modified by the effects of long-term heavy grazing. Kentucky bluegrass has become the dominant grass in this community. With continued heavy grazing pressure, rough fescue can disappear altogether (Moss and Campbell 1947). The presence of common dandelion is also indicative of past grazing pressure. Recovery of these sites to rough fescue will be slow, and could take 20-30 years (Willoughby et al 1996).

| Upper Bob Creek Community Type: Kentucky bluegrass - Rough fescue |                               |
|-------------------------------------------------------------------|-------------------------------|
| Sites: 9, 25                                                      | Range classification: primary |



|                                                                   |                             |                           |                  |
|-------------------------------------------------------------------|-----------------------------|---------------------------|------------------|
| Upper Bob Creek Community Type: Kentucky bluegrass - Rough fescue |                             |                           |                  |
| Moisture regime: submesic to mesic                                |                             | Soil drainage: well       |                  |
| Nutrient regime: mesotrophic                                      |                             | Soil subgroup: O.BL; O.EB |                  |
| Slope % (range): 12 (0-55)                                        |                             | Aspect: south             |                  |
| Upper Bob Creek Community Type: Kentucky bluegrass - Rough fescue |                             |                           |                  |
| Key and indicator species:                                        |                             |                           |                  |
| Scientific name                                                   |                             | Common name               | Canopy cover (%) |
| Graminoids:                                                       | <i>Poa pratensis</i>        | Kentucky bluegrass        | 29               |
|                                                                   | <i>Festuca scabrella</i>    | Rough fescue              | 7                |
|                                                                   | <i>Festuca idahoensis</i>   | Idaho fescue              | 6                |
|                                                                   | <i>Danthonia parryi</i>     | Parry oat grass           | 3                |
|                                                                   | <i>Carex obtusata</i>       | Blunt sedge               | 2                |
| Forbs:                                                            | <i>Achillea millefolium</i> | Yarrow                    | 6                |
|                                                                   | <i>Geum triflorum</i>       | Old man's whiskers        | 5                |
|                                                                   | <i>Taraxacum officinale</i> | Dandelion                 | 5                |
|                                                                   | <i>Potentilla gracilis</i>  | Graceful cinquefoil       | 3                |
| Shrubs:                                                           | <i>Potentilla fruticosa</i> | Shrubby cinquefoil        | 2                |

#### 4.2.5 Pine grass - Hairy wild rye / Wild strawberry

On moist open upper slopes where grassland begins to grade to forest, the dominant grasses are two species which are common understory species in woodlands, pine grass (*Calamagrostis rubescens*) and hairy wild rye (*Elymus innovatus*). This community type is transitional to forest. It was found on a steep east-facing upper slope, amid scattered lodgepole pine and spruce. The more favourable moisture regime may be due to the leeward position of the site, which would favour snow accumulation. Plant species characteristic of favourable moisture regimes were present, such as showy aster, as well as some species, such as hairy wild rye, often found on drier sites.

|                                                                              |                                 |
|------------------------------------------------------------------------------|---------------------------------|
| Upper Bob Creek Community Type: Pine grass - Hairy wild rye/ Wild strawberry |                                 |
| Sites: #7                                                                    | Range classification: Secondary |
| Moisture regime: mesic                                                       | Soil drainage: moderately well  |
| Nutrient regime: mesotrophic                                                 | Soil subgroup: O.EB             |



|                                                                              |                                                |                        |                  |
|------------------------------------------------------------------------------|------------------------------------------------|------------------------|------------------|
| Upper Bob Creek Community Type: Pine grass - Hairy wild rye/ Wild strawberry |                                                |                        |                  |
| Slope % (range): 20 (0-46)                                                   |                                                | Aspect: north; east    |                  |
| Key and indicator species:                                                   |                                                |                        |                  |
|                                                                              | Scientific name                                | Common name            | Canopy cover (%) |
| Graminoids:                                                                  | <i>Calamagrostis rubescens</i>                 | Pine grass             | 24               |
|                                                                              | <i>Elymus innovatus</i>                        | Hairy wild rye         | 3                |
|                                                                              | <i>Bromus inermis</i> var. <i>pumpellianus</i> | Awnless brome          | 1                |
| Forbs:                                                                       | <i>Fragaria virginiana</i>                     | Strawberry             | 14               |
|                                                                              | <i>Lathyrus ochroleucus</i>                    | Peavine                | 3                |
|                                                                              | <i>Lupinus sericeus</i>                        | Silky perennial lupine | 4                |
|                                                                              | <i>Aster conspicuus</i>                        | Showy aster            | 3                |
| Shrubs:                                                                      | <i>Vaccinium caespitosum</i>                   | Dwarf bilberry         | 1                |
|                                                                              | <i>Symphoricarpos occidentalis</i>             | Buckbrush              | 2                |
|                                                                              | <i>Rosa acicularis</i>                         | Prickly rose           | 1                |
|                                                                              | <i>Spiraea betulifolia</i>                     | White meadowsweet      | 2                |

## Shrubland community types

One lowland shrubland community type was sampled and it is described in detail below. However, small patches of other shrubs can also be found bordering aspen copses, as well as on the grassy slopes of smaller coulees and moister swales and nivation hollows within grassland areas. Common shrub species in these areas include prickly rose (*Rosa acicularis*), saskatoon (*Amelanchier alnifolia*), beaked willow (*Salix bebbiana*) and buckbrush or snowberry (*Symphoricarpos* sp.). Willoughby et al (1996) describe a snowberry - rose - saskatoon community occurring on south-facing slopes in transition from rough fescue grassland to forest, and O'Leary et al (1989) found a similar community on well-drained south- and southwest-facing slopes. A beaked willow / hairy wild rye community (Willoughby et al 1996) likely occurs on north-facing slopes and other areas with suitable soil moisture conditions. Shrublands provide excellent browse for wildlife (Jaques and Corbin 1981).

### 4.2.6 Bog birch - Basket willow - Myrtle-leaved willow

Extensive areas of bog birch- willow shrubland cover the wide Upper Bob Creek valley. The willow stands are heavily utilized by moose (this study; O'Leary et al 1989). This community type occurs on gently inclined and undulating recent fluvial deposits in the bottom of Bob Creek valley, especially on fluvial terraces and other areas associated with older creek channels. Soils are primarily Orthic Humic Gleysols. Soil drainage is often impeded, as these sites usually occur on seepage zones, with a water table close to the surface. The variability in drainage is expressed



in the species composition of the shrub layer. Myrtle-leaved willow predominates in the central damp areas, where soil drainage is poorest, whereas basket willow is more common on the slightly drier portions of sites. Likewise, water-loving species such as beaked sedge and wire rush are found in areas with prolonged flooding. Where temporary flooding is of shorter duration, the forbs purple avens and veiny meadow rue, and grasses bluejoint and tufted hair grass occur.

| Upper Bob Creek Community Type: Bog birch - Basket willow - Myrtle-leaved willow |                                 |                                 |                  |
|----------------------------------------------------------------------------------|---------------------------------|---------------------------------|------------------|
| Sites: 4,11, 21, 22                                                              |                                 | Range classification: Secondary |                  |
| Moisture regime: subhydryc                                                       |                                 | Soil drainage: poorly           |                  |
| Nutrient regime: permesotrophic                                                  |                                 | Soil subgroup: O.HG; T.M        |                  |
| Slope % (range): 0-6                                                             |                                 | Aspect: northerly; level        |                  |
| Key and indicator species:                                                       |                                 |                                 |                  |
| Scientific name                                                                  |                                 | Common name                     | Canopy cover (%) |
| Shrubs:                                                                          | <i>Salix petiolaris</i>         | Basket willow                   | 26               |
|                                                                                  | <i>Betula glandulosa</i>        | Bog birch                       | 11               |
|                                                                                  | <i>Salix myrtillifolia</i>      | Myrtle-leaved willow            | 9                |
|                                                                                  | <i>Potentilla fruticosa</i>     | Shrubby cinquefoil              | 2                |
|                                                                                  | <i>Salix candida</i>            | Hoary willow                    | 5                |
|                                                                                  | <i>Salix glauca</i>             | Smooth willow                   | 4                |
| Graminoids:                                                                      | <i>Juncus balticus</i>          | Wire rush                       | 5                |
|                                                                                  | <i>Carex rostrata</i>           | Beaked sedge                    | 5                |
|                                                                                  | <i>Carex praeegracilis</i>      | Graceful sedge                  | 3                |
|                                                                                  | <i>Agropyron trachycaulum</i>   | Slender wheat grass             | 2                |
|                                                                                  | <i>Calamagrostis canadensis</i> | Bluejoint                       | 1                |
|                                                                                  | <i>Deschampsia cespitosa</i>    | Tufted hair grass               | 1                |
| Forbs:                                                                           | <i>Thalictrum venulosum</i>     | Veiny meadow rue                | 3                |
|                                                                                  | <i>Epilobium angustifolium</i>  | Fireweed                        | 2                |
|                                                                                  | <i>Geum rivale</i>              | Purple avens                    | 2                |
|                                                                                  | <i>Aster laevis</i>             | Smooth aster                    | 3                |

## Deciduous forest community types

Two aspen community types were sampled in this study, representing two fairly different ecological units. One type was a transitional community on a dry site, representing an invasion of aspen into grassland, and the other was found where both moisture and nutrient conditions



were more favourable. Understory of the aspen communities reflected the differences in moisture and nutrient regimes. Bearberry was predominant at the dry nutrient-poor site, whereas prickly rose was characteristic of the moister and richer site.

#### 4.2.7 Aspen / Bearberry / Rough fescue

In upper slope areas, small groves of aspen occur on warm, protected and not too dry sites such as southeast-facing slopes protected from desiccating winds. Here aspen is growing at the edge of its ecological niche (Willoughby et al 1996). Associated species in these drier areas include bearberry, forb species such as wild white geranium (*Geranium richardsonii*), and the grasses rough fescue and hairy wild rye. It is not known to what extent aspen is invading grasslands in the Ecological Reserve. In a study in the nearby Porcupine Hills in the 1970's, aspen invasion occurred most commonly on north-facing slopes, and in some areas aspen woods were succeeding to Douglas fir (Fredrickson 1975).

The aspen/bearberry/rough fescue community type described below occurs on colluvial and morainal materials in the study area. Because of the presence of bearberry, this community type is included in the 'bearberry Aw' ecosite phase of Archibald et al (1996), defined by moisture/nutrient regimes that are submesic/poor. The presence of rough fescue, however, indicates that soils at these sites are better developed and somewhat richer than in the modal bearberry communities. This community type may represent aspen invasion into a rough fescue - sedge/bearberry community type, described by Willoughby et al (1996) and characteristic of dry grass meadows on hilltops in the Montane.

| Upper Bob Creek Community Type: Aspen/Bearberry/Rough fescue |                                |                                 |                  |
|--------------------------------------------------------------|--------------------------------|---------------------------------|------------------|
| Sites: #46                                                   |                                | Range classification: secondary |                  |
| Moisture regime: submesic                                    |                                | Soil drainage: well             |                  |
| Nutrient regime: mesotrophic                                 |                                | Soil subgroup: O.EB; O.DG       |                  |
| Slope% (range): 5 (0-10)                                     |                                | Aspect: hilltop; south          |                  |
| Key and indicator species:                                   |                                |                                 |                  |
| Scientific name                                              |                                | Common name                     | Canopy cover (%) |
| Trees:                                                       | <i>Populus tremuloides</i>     | Aspen                           | 12               |
| Shrubs:                                                      | <i>Arctostaphylos uva-ursi</i> | Common bearberry                | 13               |
|                                                              | <i>Rosa acicularis</i>         | Prickly rose                    | 1                |
|                                                              | <i>Potentilla fruticosa</i>    | Shrubby cinquefoil              | 2                |



|             |                                |                 |    |
|-------------|--------------------------------|-----------------|----|
| Forbs:      | <i>Aster ciliolatus</i>        | Lindley's aster | 10 |
|             | <i>Fragaria virginiana</i>     | Wild strawberry | 5  |
|             | <i>Achillea millefolium</i>    | Yarrow          | 4  |
|             | <i>Taraxacum officinale</i>    | Dandelion       | 5  |
| Graminoids: | <i>Festuca scabrella</i>       | Rough fescue    | 15 |
|             | <i>Festuca idahoensis</i>      | Idaho fescue    | 12 |
|             | <i>Calamagrostis rubescens</i> | Pine grass      | 11 |

#### 4.2.8 Aspen / Rose / Pine grass

This community type was found at 4 sites during the study, located on a variety of aspects except south-facing slopes, usually on lower slope positions. It occurred most commonly on moderately well to well-drained Orthic Black Chernozems developed on morainal parent materials. These lower slope aspen stands occur on mesic, well-drained sites with medium to rich nutrient regimes. The most common understory shrubs in these areas are prickly rose, saskatoon and snowberry (*Symphoricarpos albus*). Characteristic forbs include showy aster (*Aster conspicuus*), cream-coloured vetchling (*Lathyrus ochroleucus*) and wild strawberry (*Fragaria virginiana*), while the most common grass species is pine grass. A similar aspen/ pine grass community type was described by Archibald et al (1996), and included in their 'thimbleberry / pine grass' (mesic / rich) Aw (aspen) ecosite phase. Succession in this community is towards white spruce (*Picea glauca*).

|                                                       |                              |                                 |                  |
|-------------------------------------------------------|------------------------------|---------------------------------|------------------|
| Upper Bob Creek Community Type: Aspen/Rose/Pine grass |                              |                                 |                  |
| Sites: 31, 34, 38, 47                                 |                              | Range classification: Secondary |                  |
| Moisture regime: mesic                                |                              | Soil drainage: well             |                  |
| Nutrient regime: permesotrophic to mesotrophic        |                              | Soil subgroup: O.BL; O.EB       |                  |
| Slope(range): 5-20%                                   |                              | Aspect: north, west, east       |                  |
| Key and indicator species:                            |                              |                                 |                  |
| Scientific name                                       |                              | Common name                     | Canopy cover (%) |
| Trees:                                                | <i>Populus tremuloides</i>   | Aspen                           | 41               |
|                                                       | <i>Populus balsamifera</i>   | Balsam poplar                   | 3                |
| Shrubs:                                               | <i>Rosa acicularis</i>       | Prickly rose                    | 8                |
|                                                       | <i>Amelanchier alnifolia</i> | Saskatoon                       | 2                |
|                                                       | <i>Spiraea betulifolia</i>   | White meadowsweet               | 1                |



|             |                                |                          |    |
|-------------|--------------------------------|--------------------------|----|
| Forbs:      | <i>Fragaria virginiana</i>     | Wild strawberry          | 8  |
|             | <i>Aster ciliolatus</i>        | Lindley's aster          | 7  |
|             | <i>Aster conspicuus</i>        | Showy aster              | 7  |
|             | <i>Lathyrus ochroleucus</i>    | Cream-coloured vetchling | 5  |
|             | <i>Galium boreale</i>          | Northern bedstraw        | 2  |
|             | <i>Vicia americana</i>         | Vetch                    | 3  |
|             |                                |                          |    |
| Graminoids: | <i>Calamagrostis rubescens</i> | Pine grass               | 16 |
|             | <i>Elymus innovatus</i>        | Hairy wild rye           | 6  |

## Mixed forest community types

Mixed forest community types represent mid-successional stages towards conifer-dominated types. Two mixed forest community types were sampled, an aspen - spruce community and a spruce - balsam community. However, other mixed forest communities are no doubt present in the study area, especially on northeast or northwest facing slopes. Mixed spruce-balsam poplar forests occur on the most nutrient rich sites, usually in coulee bottoms or adjacent to the Bob Creek watercourse. Mixed stands of aspen and Douglas fir are often present on mid-slope positions with northwest or northeast exposures. Several mixed forest types are discussed by Willoughby et al (1996) including aspen - Douglas fir / white meadowsweet, aspen -lodgepole pine / buffaloberry / hairy wild rye, and aspen - Douglas fir / bearberry. O'Leary et al (1989) found an aspen - white spruce / rose / reed grass type on mid to lower slopes (aspects east, northeast and northwest). Archibald et al (1996) describe an aspen - spruce - lodgepole pine - Douglas fir mixed forest on mesic sites with mesotrophic nutrient regimes.

The two mixed forest community types sampled in this study are described below.

### 4.2.9 Aspen - White spruce/ Blueberry

This community type occurred on a north-facing lower slope area underlain by morainal and colluvial veneers over bedrock. Soils were moderately well to well-drained Orthic Dark Grey Chernozems, with a mesic moisture regime. Nutrient regime was mesotrophic, somewhat less rich than the aspen / rose / pine grass community type described above. The presence of grouseberry may indicate cooler conditions at this site, since this species is more frequent at higher elevations, in a subalpine environment. Succession is to a white spruce/moss community (Willoughby et al 1996). This community type would be classed within a 'white meadowsweet Sw' (white spruce) ecosite phase (mesic moisture and medium nutrient regimes) (Archibald et al 1996).

|                                                                |                               |
|----------------------------------------------------------------|-------------------------------|
| Upper Bob Creek Community Type: Aspen- White spruce/ Blueberry |                               |
| Sites: #37                                                     | Range classification: non-use |
| Moisture regime: mesic                                         | Soil drainage: well           |



|                                                                |                                    |                          |                  |
|----------------------------------------------------------------|------------------------------------|--------------------------|------------------|
| Upper Bob Creek Community Type: Aspen- White spruce/ Blueberry |                                    |                          |                  |
| Nutrient regime: mesotrophic                                   |                                    | Soil subgroup: O.DG      |                  |
| Slope: 2%                                                      |                                    | Aspect: north            |                  |
| Key and indicator species:                                     |                                    |                          |                  |
| Scientific name                                                |                                    | Common name              | Canopy cover (%) |
| Trees:                                                         | <i>Populus tremuloides</i>         | Aspen                    | 33               |
|                                                                | <i>Picea glauca</i>                | White spruce             | 36               |
| Shrubs:                                                        | <i>Vaccinium membranaceum</i>      | Tall bilberry            | 6                |
|                                                                | <i>Vaccinium scoparium</i>         | Grouse-berry             | 7                |
|                                                                | <i>Rosa acicularis</i>             | Prickly rose             | 1                |
|                                                                | <i>Symphoricarpos occidentalis</i> | Buckbrush                | 4                |
| Forbs:                                                         | <i>Fragaria virginiana</i>         | Wild strawberry          | 8                |
|                                                                | <i>Lathyrus ochroleucus</i>        | Cream-coloured vetchling | 2                |
|                                                                | <i>Thalictrum occidentale</i>      | Western meadow rue       | 4                |
|                                                                | <i>Pyrola asarifolia</i>           | Common pink wintergreen  | 2                |
|                                                                | <i>Vicia americana</i>             | American vetch           | 1                |
|                                                                | <i>Galium boreale</i>              | Northern bedstraw        | 1                |
| Graminoids:                                                    | <i>Calamagrostis rubescens</i>     | Pine grass               | 3                |
|                                                                | <i>Melica smithii</i>              | Melic grass              | 12               |

#### 4.2.10 Spruce - Balsam poplar / Snowberry

This community type occurred in a coulee bottom at a fairly high elevation (1569 m). In the coulees, Orthic Eutric Brunisolic soils are interspersed with Orthic Humic Gleysols on fluvial veneers over bedrock (Leskiw 1993). The characteristic subhygric moisture regime has allowed these sites to escape recent fire, allowing succession towards the white spruce climax. Archibald et al (1996) describe a similar balsam poplar / snowberry community, which is at an earlier successional stage. It grows on sites with subhygric/rich moisture/nutrient regimes.

|                                                                         |                                 |
|-------------------------------------------------------------------------|---------------------------------|
| Upper Bob Creek Community Type: White spruce - Balsam poplar/ Snowberry |                                 |
| Sites: #45                                                              | Range classification: Secondary |
| Moisture regime: subhygric                                              | Soil drainage: moderately well  |
| Nutrient regime: mesotrophic to permesotrophic                          | Soil subgroup: O.EB             |
| Slope: level                                                            | Aspect: level                   |
| Key and indicator species:                                              |                                 |



| Scientific name |                                 | Common name              | Canopy cover (%) |
|-----------------|---------------------------------|--------------------------|------------------|
| Trees:          | <i>Populus balsamifera</i>      | Balsam poplar            | 30               |
|                 | <i>Picea glauca</i>             | White spruce             | 25               |
|                 | <i>Populus tremuloides</i>      | Aspen                    | 5                |
|                 |                                 |                          |                  |
| Shrubs:         | <i>Symphoricarpos albus</i>     | Snowberry                | 6                |
|                 | <i>Rosa acicularis</i>          | Prickly rose             | 3                |
|                 | <i>Rubus idaeus</i>             | Raspberry                | 5                |
|                 |                                 |                          |                  |
| Forbs:          | <i>Lathyrus ochroleucus</i>     | Cream-coloured vetchling | 3                |
|                 | <i>Geranium richardsonii</i>    | Wild white geranium      | 5                |
|                 | <i>Aster ciliolatus</i>         | Lindley's aster          | 6                |
|                 | <i>Viola canadensis</i>         | Western Canada violet    | 3                |
|                 | <i>Fragaria virginiana</i>      | Wild strawberry          | 4                |
|                 | <i>Epilobium angustifolium</i>  | Fireweed                 | 4                |
|                 |                                 |                          |                  |
| Graminoids:     | <i>Calamagrostis canadensis</i> | Bluejoint                | 7                |

### Conifer forest community types

A variety of closed and open coniferous forest types, dominated by white spruce, Douglas fir, lodgepole pine and limber pine, occur in the Reserve. White spruce occurs on the most nutrient rich sites in the study area, usually in coulee bottoms or adjacent to the Bob Creek watercourse. Douglas fir is found on relatively dry, mid-slope to upper slope locations of varying aspects, which have escaped fire. Lodgepole pine is a fire successional species, and thus its presence is indicative of a local history of fire. It generally occurs on relatively dry sites at mid-slope or higher elevations in the study area. Limber pine occurs on dry (subxeric) upper slope sites, such as at the tops of ridges, often on poorly developed soils (Orthic Regosols). This species is well-adapted to low soil moisture and nutrient levels (Kuchar 1973), but occurs only in small patches in the study area.

Because of a relatively high frequency of fire, closed-canopied lodgepole pine forests are the modal forest for the Montane subregion. In the absence of fire, succession is towards Douglas fir on the drier sites and white spruce on the more moist sites (Strong and Leggat 1992).

Site moisture and nutrient conditions are important factors controlling forest understory composition and potential climax forest. For example, the driest and most nutrient-poor lodgepole pine stands have an understory layer dominated by bearberry, grading to white meadowsweet (*Spiraea betulifolia*) in the more mesotrophic sites, with thimbleberry (*Rubus parviflorus*) the dominant shrub on the richest and moistest sites (Archibald et al 1996).

Forested community types found in the Upper Bob Creek Ecological Reserve are described below. No limber pine communities were sampled in the study area, but they are likely present on very dry, nutrient-poor sites on exposed ridges.



#### 4.2.11 Douglas fir /Hairy wild rye

This community type was found at 4 sites in the study area. It occurred on steep, dry sites at a range of elevations (1460 to 1708 m). At these sites, shallow Eutric Brunisolic soils had developed on colluvial and residual (eroded bedrock) parent materials. Aspect was west-facing or east-facing at sites examined in this study, though the type has been found on a variety of aspects (Willoughby et al 1996). The type is one of a number of coniferous community types occurring where site moisture and nutrient regimes are, respectively, submesic and medium (Archibald et al 1996). In the Porcupine Hills, a Montane area to the east, Jaques (1979) found that at lower elevations, below 1400 m, Douglas fir occupied north- or east- facing aspects, while at higher elevations it was found on all aspects.

| Upper Bob Creek Community Type: Douglas fir/Hairy wild rye |                                |                               |                  |
|------------------------------------------------------------|--------------------------------|-------------------------------|------------------|
| Sites: 33,35,36,39                                         |                                | Range classification: non-use |                  |
| Moisture regime: submesic to mesic                         |                                | Soil drainage: well           |                  |
| Nutrient regime: mesotrophic                               |                                | Soil subgroup: O.EB           |                  |
| Slope: 15-45%                                              |                                | Aspect: west, east            |                  |
| Key and indicator species:                                 |                                |                               |                  |
| Scientific name                                            |                                | Common name                   | Canopy cover (%) |
| Trees:                                                     | <i>Pseudotsuga menziesii</i>   | Douglas fir                   | 60               |
|                                                            | <i>Pinus contorta</i>          | Lodgepole pine                | 1                |
| Shrubs:                                                    | <i>Rosa acicularis</i>         | Prickly rose                  | 3                |
|                                                            | <i>Spiraea betulifera</i>      | White meadowsweet             | 6                |
|                                                            | <i>Juniperus communis</i>      | Ground juniper                | 2                |
| Forbs:                                                     | <i>Aster conspicuus</i>        | Showy aster                   | 5                |
|                                                            | <i>Arnica cordifolia</i>       | Heart-leaved arnica           | 5                |
|                                                            | <i>Fragaria virginiana</i>     | Wild strawberry               | 4                |
|                                                            | <i>Thalictrum venulosum</i>    | Veiny meadow rue              | 2                |
|                                                            | <i>Lathyrus ochroleucus</i>    | Cream-coloured vetchling      | 1                |
|                                                            | <i>Galium boreale</i>          | Northern bedstraw             | 1                |
| Graminoids:                                                | <i>Elymus innovatus</i>        | Hairy wild rye                | 6                |
|                                                            | <i>Calamagrostis rubescens</i> | Pine grass                    | 3                |

#### 4.2.12 Lodgepole pine/ White meadowsweet

This community type is included in a group of community types growing on sites with mesic moisture regimes and medium nutrient regimes (Archibald et al 1996), which includes types



4.2.13 and 4.2.14 discussed below. The dominant overstory species and the cover of white meadowsweet vary with the successional status of the site (Willoughby et al 1996). The lodgepole pine/white meadowsweet community type was found on an east-facing slope at an elevation of 1569 m, and was dominated by pine in the overstory. It occurred on Orthic Eutric Brunisols, which had developed on till (Leskiw 1993). A history of fire is responsible for maintaining the pine overstory. Succession in the lodgepole pine/meadowsweet community type is to white spruce or Douglas fir or both (Archibald et al 1996).

| Upper Bob Creek Community Type: Lodgepole pine/ white meadowsweet |                                                                                                                    |                                                                         |                  |
|-------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------|------------------|
| Sites: #40                                                        |                                                                                                                    | Range classification: non-use                                           |                  |
| Moisture regime: mesic                                            |                                                                                                                    | Soil drainage: well                                                     |                  |
| Nutrient regime: medium                                           |                                                                                                                    | Soil subgroup: O.EB                                                     |                  |
| Slope: 10-53%                                                     |                                                                                                                    | Aspect: east(this study); south to southwest<br>(Willoughby et al 1996) |                  |
| Key and indicator species:                                        |                                                                                                                    |                                                                         |                  |
|                                                                   | Scientific name                                                                                                    | Common name                                                             | Canopy cover (%) |
| Trees                                                             | <i>Pinus contorta</i><br><i>Populus tremuloides</i>                                                                | Lodgepole pine<br>Aspen                                                 | 66<br>1          |
| Shrubs:                                                           | <i>Spiraea betulifolia</i><br><i>Shepherdia canadensis</i><br><i>Rosa acicularis</i>                               | White meadowsweet<br>Canada buffaloberry<br>Prickly rose                | 18<br>1<br>3     |
| Forbs:                                                            | <i>Fragaria virginiana</i><br><i>Arnica cordifolia</i><br><i>Aster conspicuus</i><br><i>Vaccinium membranaceum</i> | Wild strawberry<br>Heart-leaved arnica<br>Showy aster<br>Tall bilberry  | 4<br>4<br>3<br>9 |
| Graminoids:                                                       | <i>Elymus innovatus</i><br><i>Calamagrostis rubescens</i><br><i>Melica smithii</i>                                 | Hairy wild rye<br>Pine grass<br>Melic grass                             | 1<br>10<br>2     |

#### 4.2.13 White spruce- Douglas fir/ White meadowsweet

This type is characterized by white spruce and Douglas fir in the overstory. It occurs on sites similar to the lodgepole pine/meadowsweet community type described above, but is successional more mature. Moisture conditions are slightly more favourable in this community type.



|                                                                             |                                                                                                             |                                                                             |                   |
|-----------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|-------------------|
| Upper Bob Creek Community Type: White spruce-Douglas fir/ White meadowsweet |                                                                                                             |                                                                             |                   |
| Sites: #42, 43                                                              |                                                                                                             | Range classification: non-use (secondary)                                   |                   |
| Moisture regime: mesic                                                      |                                                                                                             | Soil drainage: moderately well to well                                      |                   |
| Nutrient regime: mesotrophic                                                |                                                                                                             | Soil subgroup: O.EB; O.DG                                                   |                   |
| Slope: 20-40%                                                               |                                                                                                             | Aspect: north, west                                                         |                   |
| Key and indicator species:                                                  |                                                                                                             |                                                                             |                   |
| Scientific name                                                             |                                                                                                             | Common name                                                                 | Canopy cover (%)  |
| Trees:                                                                      | <i>Picea glauca</i><br><i>Pseudotsuga menziesii</i><br><i>Pinus contorta</i>                                | White spruce<br>Douglas fir<br>Lodgepole pine                               | 33<br>20<br>7     |
| Shrubs:                                                                     | <i>Spiraea betulifolia</i><br><i>Rubus parviflorus</i><br><i>Rosa acicularis</i>                            | White meadowsweet<br>Thimbleberry<br>Prickly rose                           | 6<br>5<br>1       |
| Forbs:                                                                      | <i>Thalictrum venulosum</i><br><i>Arnica cordifolia</i><br><i>Aster conspicuus</i><br><i>Galium boreale</i> | Veiny meadow rue<br>Heart-leaved arnica<br>Showy aster<br>Northern bedstraw | 10<br>6<br>5<br>1 |
| Graminoids:                                                                 | <i>Calamagrostis rubescens</i><br><i>Elymus innovatus</i>                                                   | Pine grass<br>Hairy wild rye                                                | 2<br>1            |

#### 4.2.14 White spruce/moss

The white spruce/moss community type is the predicted climax community type on sites occupied by the lodgepole pine/ white meadowsweet and the Douglas fir - white spruce/ white meadowsweet community types. In the study area it was found primarily on north-facing slopes or adjacent to drainages. The more favourable soil moisture conditions at these sites have likely helped to protect these sites from fire. The most common underlying parent materials are colluvial, and soils are Orthic Eutric Brunisols (Leskiw 1993). White spruce is the undisputed dominant in the overstory, although a few aspen occur. As a result of the dense spruce overstory, cover of grasses and forbs is low in this community type.

|                                                    |                                        |
|----------------------------------------------------|----------------------------------------|
| Upper Bob Creek Community Type: White spruce/ moss |                                        |
| Sites: #30, 32, 41                                 | Range classification: non-use          |
| Moisture regime: mesic                             | Soil drainage: moderately well to well |
| Nutrient regime: mesotrophic                       | Soil subgroup: O.EB                    |



|                                                    |                             |                       |                         |
|----------------------------------------------------|-----------------------------|-----------------------|-------------------------|
| Upper Bob Creek Community Type: White spruce/ moss |                             |                       |                         |
| Slope % (range): 5-25                              |                             | Aspect: north         |                         |
| Key and indicator species:                         |                             |                       |                         |
|                                                    | <b>Scientific name</b>      | <b>Common name</b>    | <b>Canopy cover (%)</b> |
| Trees:                                             | <i>Picea glauca</i>         | White spruce          | 64                      |
|                                                    | <i>Populus tremuloides</i>  | Douglas fir           | 2                       |
| Shrubs:                                            | <i>Rosa acicularis</i>      | Prickly rose          | 2                       |
|                                                    | <i>Linnaea borealis</i>     | Twin-flower           | 8                       |
| Forbs:                                             | <i>Arnica cordifolia</i>    | Heart-leaved arnica   | 6                       |
|                                                    | <i>Thalictrum venulosum</i> | Veiny meadowrue       | 1                       |
|                                                    | <i>Aster conspicuus</i>     | Showy aster           | 1                       |
|                                                    | <i>Pyrola asarifolia</i>    | Pink wintergreen      | 1                       |
|                                                    | <i>Orthilia secunda</i>     | One-sided wintergreen | 1                       |
| Graminoids:                                        | <i>Carex spp.</i>           | Sedge                 | 2                       |
|                                                    | <i>Elymus innovatus</i>     | Hairy wild rye        | 1                       |
| Moss:                                              | <i>Hylocomium splendens</i> | Stair-step moss       | 49                      |
|                                                    | <i>Pleurozium schreberi</i> | Schreber's moss       | 5                       |

#### 4.2.15 White spruce/ Thimbleberry

This community type occurs on ecosites with moisture/nutrient regimes that are respectively mesic/rich (Archibald et al 1996). The presence of thimbleberry is indicative of the better than average nutrient conditions, which may be a result of local nutrient-rich seepage. This community type was found only at one site, on Orthic Black Chernozemic soils which had developed on morainal parent materials on a lower slope west of Bob Creek valley. The dominant species in the overstory is white spruce, which represents the climax overstory species at this site.

|                                                            |                                |
|------------------------------------------------------------|--------------------------------|
| Upper Bob Creek Community Type: White spruce/ Thimbleberry |                                |
| Sites: #44                                                 | Range classification: non-use  |
| Moisture regime: mesic to subhygric                        | Soil drainage: moderately well |
| Nutrient regime: mesotrophic to permesotrophic             | Soil subgroup: O.BL; O.EB      |
| Slope: 15%                                                 | Aspect: north                  |
| Key and indicator species:                                 |                                |



| Scientific name |                          | Common name         | Canopy cover (%) |
|-----------------|--------------------------|---------------------|------------------|
| Trees:          | <i>Picea glauca</i>      | White spruce        | 75               |
| Shrubs:         | <i>Rubus parviflorus</i> | Thimbleberry        | 11               |
|                 | <i>Cornus canadensis</i> | Bunchberry          | 12               |
|                 | <i>Linnaea borealis</i>  | Twinflower          | 4                |
|                 | <i>Rosa acicularis</i>   | Prickly rose        | 1                |
| Forbs           | <i>Arnica cordifolia</i> | Heart-leaved arnica | 3                |
|                 | <i>Aster conspicuus</i>  | Showy aster         | 2                |
|                 | <i>Angelica dawsonii</i> | Yellow angelica     | 1                |
| Graminoids:     | <i>Melica smithii</i>    | Melic grass         | 1                |

#### 4.2.16 Limber pine - Douglas fir/Juniper - Bearberry (Willoughby et al 1996)

This community type was not sampled during this study. However, limber pine-dominated community types likely form an edaphic climax on shallow-soiled, steep ridges of high elevation, where bedrock exposures are frequent, such as in the northwest part of the study area. These are areas exposed to high winds. A limber pine-Douglas fir/ Juniper- Bearberry community type was described by Willoughby et al (1996). The presence of bearberry and juniper is indicative of dry, nutrient-poor soil conditions.

#### 4.3 Significant Plant Species

At least 10 rare or uncommon plant species have been recorded at Upper Bob Creek Ecological Reserve in this study or by previous workers (Table 1). Details of their status, habitat requirements (Moss 1983; Wallis et al 1987) and details of occurrence are given below. Moist montane meadows appear to offer habitat for the greatest potential number of rare plants. A number of other species may occur in the study area, but their presence is not confirmed (Appendix D).

Sources used to determine the status of plant species found in the Ecological Reserve include the following: Packer and Bradley (1984), Wallis et al (1986) and (1987), Fairbarns et al (1987) and Alberta Natural Heritage Information Centre (1996). Species rare in Alberta were defined by Packer and Bradley (1984) as those with known collections from five or fewer localities. The status of these and several other species was further assessed by Wallis et al (1986) and (1987), and Fairbarns et al (1987). These lists generally include species which are rare due to limited geographical distribution, and those which occur in small numbers wherever they grow. A list of Plant Species of Concern has also been recently collated by the Alberta Natural Heritage Information Centre (ANHIC 1996), which updates the information provided in other references. This list identifies species of special concern or high priority because they are rare, endemic, disjunct or in peril. Species are ranked based primarily on number of occurrences, with



consideration also given to distribution and current threats to populations. These ranking classes are summarized in Appendix E.

Significant plant species at Upper Bob Creek Ecological Reserve:

1) *Angelica dawsonii* S. Watts (Yellow angelica)

**Status:** restricted southwest Alberta range, but not uncommon (Wallis et al 1986);

**Habitat:** moist meadows and woods along streams

**Occurrence:** recorded by Brown et al (1986); and this study (sites 32-34, 37, 41, 42, 44, 45)

2) *Balsamorhiza sagittata* (Pursh) Nutt. (Balsamroot)

**Status:** restricted southwest Alberta distribution (Moss 1983); representative Montane species, but not uncommon

**Habitat:** montane grassland and open woods; forest edges;

**Occurrence:** found at several sites in the Reserve (this study)

3) *Camassia quamash* (Pursh) Greene (Blue camas)

**Status:** species of special concern; S2 on Alberta tracking list; uncommon in Alberta (7 mapped locations in the province); found only in the southwest part of the province; moderately threatened due to habitat loss (ANHIC 1996)

**Habitat:** wet meadows

**Occurrence:** recorded in the study area (ANHIC database, 1996); also Brown et al (1986)

4) *Carex scopulorum* Holm. (Holm's Rocky Mountain sedge)

**Status:** uncommon in Alberta (in foothills areas; Moss 1983);

**Habitat:** moist open areas

**Occurrence:** recorded in this study, and by Brown et al (1986)

5) *Castilleja cusickii* Greenm. (Yellow paintbrush)

**Status:** species of special concern ranked S2 on ANHIC tracking list; rare in Alberta (Packer and Bradley 1984); recommended classification as 'endangered' (Wallis et al 1986);

**Habitat:** moist meadows and grassland

**Occurrence:** recorded this study (site #13)

6) *Conimitella williamsii* (D.C. Eat.) Rydb. (Conimitella)

**Status:** species of special concern - S2 on ANHIC tracking list; only 6-20 records (ANHIC 1996)

**Habitat:** open montane, grassland slopes

**Occurrence:** recorded in this study (sites #7, 10); and by Brown et al (1986)

7) *Corallorhiza striata* Lindl. (Striped coralroot)

**Status:** uncommon and scattered in Alberta (Moss 1983); habitat moderately threatened (J. Rintoul, Alberta Natural Areas Program, pers. comm.)

**Habitat:** woods

**Occurrence:** recorded this study



Table 1. Rare or uncommon plants recorded at Upper Bob Creek Ecological Reserve.

| Species                                               | Common name                    | Ranking<br>(ANHIC) |
|-------------------------------------------------------|--------------------------------|--------------------|
| <b>Species of concern <sup>a</sup> :</b>              |                                |                    |
| <i>Castilleja cusickii</i>                            | Yellow paintbrush              | S2                 |
| <i>Camassia quamash</i>                               | Blue camas                     | S2                 |
| <i>Conimitella williamsii</i>                         | Conimitella                    | S2                 |
| <i>Poa nevadensis</i>                                 | Nevada bluegrass               | SU                 |
| <i>Senecio foetidus</i> var.<br><i>hydrophiloides</i> | Marsh butterweed               | S3                 |
| <b>Other significant species:</b>                     |                                |                    |
| <i>Angelica dawsonii</i>                              | Yellow angelica                | ---                |
| <i>Balsamorhiza sagittata</i>                         | Balsamroot                     | ---                |
| <i>Carex scopulorum</i>                               | Holm's rocky mountain<br>sedge | ---                |
| <i>Corallorhiza striata</i>                           | Striped coralroot              | ---                |
| <i>Osmorhiza occidentalis</i>                         | Western sweet cicely           | ---                |

<sup>a</sup> Alberta Natural Heritage Information Centre: Tracking list of plant species of special concern.



8) *Osmorhiza occidentalis* (Nutt.) Torr. (Western sweet cicely)

**Status:** only 5 records in Alberta (Moss 1983); restricted range in Alberta (Wallis et al 1986; )

**Habitat:** montane woods

**Occurrence:** recorded this study (site 10), and by Brown et al (1986)

9) *Poa nevadensis* Vasey ex Scribn. (Nevada bluegrass)

**Status:** identified as a priority species, but status not confirmed (ANHIC 1996); rare in Alberta (Wallis et al 1987; Packer and Bradley 1984)

**Habitat:** moist meadows

**Occurrence:** recorded this study

10) *Senecio foetidus* Howell var. *hydrophiloides* (Marsh butterweed)

**Status:** identified as rare in Alberta (Packer and Bradley 1984; Wallis et al 1987); ranked as S3 - locally abundant, but has a limited distribution (ANHIC 1996)

**Habitat:** moist montane meadows

**Occurrence:** recorded this study (sites 34, 37, 38, 40, 41, 45), and by Brown et al (1986)



## 5.0 LITERATURE CITED

- Achuff, P.L. 1994. Natural regions, subregions and natural history themes of Alberta. Prep. for Alberta Environmental Protection. Edmonton. 72 pp.
- Alberta Environmental Protection. 1994. Ecological land survey site description manual. Canada-Alberta Partnership Agreement in Forestry. Edmonton, Alberta. 165 pp.
- Alberta Environmental Protection. 1993. Alberta plants and fungi - master species list and species group checklists. Pub. No. Ref. 75. Edmonton, Alberta.
- Alberta Forest Service. 1990. Range Survey Manual. 3rd Ed. Alberta Forestry, Lands and Wildlife, Edmonton.
- ANHIC (Alberta Natural Heritage Information Centre). 1996. Plant species of special concern. Alberta Environmental Protection. Edmonton.
- Archibald, J.H., G.D. Klappstein and I.G.W. Corns. 1996. Field guide to ecosites of southwestern Alberta. Spec. Rep.#8. Nor. Forestry Centre, Can. For. Serv. Edmonton.
- Brown, G., E. Gasser and P. McIsaac. 1986. Biophysical features and land use, Upper Bob Creek Candidate Ecological Reserve. Prep. for Alberta Recreation and Parks. 20 pp. plus appendices.
- Canada Soil Survey Committee. 1978. The Canadian System of Soil Classification. Canada Dept. of Agric. Public. 1646.
- Environment Canada. 1982. Canadian climate normals - temperature and precipitation. Atmospheric Environment Service. Environment Canada.
- Fairbairns, M., V. Loewen and C. Bradley. 1987. The rare vascular flora of Alberta. Vol. 1. A summary of the taxa occurring in the Rocky Mountain Natural Region. Alberta Forestry, Lands and Wildlife. Natural Areas Techn. Report No. 29. 27 pp.
- Fredrickson, D. 1975. Aspen communities and invasion of grassland in the Porcupine Hills, southwestern Alberta. M.Sc. Thesis, Dept. of Geography, Univ. of Calgary, Alberta.
- Gauch, H.G. 1982. Multivariate analysis in community ecology. Cambridge Univ. Press, Cambridge. 298 pp.
- Green, R. 1972. Geological map of Alberta. Research Council of Alberta. Edmonton.
- Jaques, D.R. 1979. Biophysical classification, effects of grazing, and biomass estimation using LANDSAT imagery on native fescue grasslands of southwestern Alberta. Prep. for Univ. of Calgary, Alberta.



- Jaques, D. and J. Corbin. 1981. Shrub communities in the Elbow and Sheep River drainages, Kananaskis Country, Alberta: their floristic composition, distribution and browse availability/use by animals. Univ. of Calgary, Alberta.
- Johnston, A. 1961. Comparison of lightly grazed and ungrazed ranged in the fescue grassland of southwestern Alberta. *Can. J. Plant Sci.* 41: 615 - 622.
- Kuchar, P. 1973. Habitat types of Waterton Lakes National Park. Alberta Energy and Natural Resources. 301 pp.
- Leskiw, L.A. 1993. Physical land classification of the Upper Bob Creek Ecological Reserve. Prep. for Alberta Environmental Protection. Edmonton. 11 pp.
- Moss, E.H. 1983. Flora of Alberta. 2nd. Ed. revised by J. Packer. Univ. Toronto Press. 687 pp.
- Moss, E.H. and J.A. Campbell. 1947. The fescue grassland of Alberta. *Can. J. Res.* 25: 209 - 227.
- O'Leary, D., J. Bentz and E. Anderson. 1989. Bob Creek winter range habitat study. Alberta Forestry, Lands and Wildlife. Edmonton.
- Packer, J. and C. Bradley. 1984. A checklist of the rare vascular plants of Alberta. Prov. Museum of Alberta, Natural History Occ. Paper No. 5. Alberta Culture, Edmonton.
- Pettapiece, W.W. 1986. Physiographic subdivisions of Alberta. Land Resource Research Centre, Research Branch, Alberta Agriculture.
- Strong, W.L. 1979. Ecological land classification and evaluation: Livingstone - Porcupine. Resource Evaluation Branch. Alberta Energy and Natural Resources.
- Strong, W.L. and K.R. Leggat. 1992. Ecoregions of Alberta. Alberta Forestry, Lands and Wildlife. Public. No. T/245. Edmonton.
- Wallis, C. C. Bradley, M. Fairbarns, J. Packer and C. Wershler. 1986. Pilot rare plant monitoring program in the Oldman Regional Plan area of southwestern Alberta. Alberta Forestry, Lands and Wildlife Publ. No. T/148. Edmonton. 203 pp.
- Wallis, C., C. Bradley, M. Fairbarns and V. Loewen. 1987. The rare flora of Alberta. Vol. 3. Species summary sheets. Alberta Forestry, Lands and Wildlife. Rep. T/155.
- Willoughby, M., M. Alexander and K. Sundquist. 1996. Range plant community types and carrying capacity for the montane subregion, second approximation. Alberta Environ. Protection, Edmonton.
- Wroe, R.A., S. Smoliak, B.W. Adams, W.D. Willms and M.L. Anderson. 1988. Guide to range condition and stocking rates for Alberta grasslands. Alberta Public Lands. Edmonton.



APPENDIX A. Map 1 polygon database. - Upper Bob Creek Ecological Reserve.

| MAP<br>POL. | PM<br>CODE* | SOIL1  | SOIL2 | MOISTURE/NUTRIENT<br>REGIMES | VEG1 | VEG2 | VEGETATION DESCRIPTION **               |
|-------------|-------------|--------|-------|------------------------------|------|------|-----------------------------------------|
| 1           | C1          | O.EB   |       | submesic/medium              | GL   |      | fescue grassland                        |
| 2           | C1          | O.EB   |       | submesic/medium              | c1   |      | Canada buffaloberry/HWR Fd              |
| 3           | F1          | O.HG   |       | mesic/rich                   | e2   |      | thimbleberry/pine grass Aw              |
| 4           | C1          | O.EB   |       | submesic/poor                | b2   |      | bearberry Aw                            |
| 5           | C1          | O.EBsh |       | submesic/medium              | c1   | c3   | Canada buffaloberry/HWR Fd; also Aw     |
| 6           | F1          | O.HG   |       | mesic/rich                   | e2   |      | thimbleberry/pine grass Aw              |
| 7           | C1          | O.EB   |       | subxeric/medium              | GL   |      | dry grassland                           |
| 8           | C1          | O.EB   |       | submesic/medium              | c1   |      | Canada buffaloberry/HWR Fd              |
| 9           | C1          | O.EB   |       | mesic/medium                 | d3   |      | white meadowsweet Sw                    |
| 10          | C1          | O.EB   |       | submesic/medium              | GL   |      | grassland                               |
| 11          | C1          | O.EB   |       | submesic/medium              | c2   |      | Canada buffaloberry/HWR PI              |
| 12          | F1          | O.HG   |       | mesic/rich                   | e2   |      | thimbleberry/pine grass Aw              |
| 13          | C1          | O.EB   |       | submesic/medium              | GL   |      | grassland                               |
| 14          | C1          | O.EB   |       | submesic/poor                | b2   |      | bearberry Aw                            |
| 15          | C1          | O.EB   |       | submesic/medium              | c1   |      | Canada buffaloberry/HWR Fd              |
| 16          | C1          | O.EB   |       | mesic/medium                 | d1   |      | white meadowsweet Fd                    |
| 17          | C1          | O.EB   |       | submesic/medium              | c2   | c1   | Canada buffaloberry/HWR PI; Fd          |
| 18          | C1          | O.EB   |       | submesic/medium              | c1   |      | Canada buffaloberry/HWR Fd              |
| 19          | C1          | O.EB   |       | mesic/medium                 | d3   |      | white meadowsweet Sw                    |
| 20          | M1          | O.BL   | O.HG  | mesic/rich                   | e2   | e3   | thimbleberry/pine grass Aw; Sw          |
| 21          | C1          | O.BL   |       | submesic/medium              | GL   | c3   | grassland with patches of poplar        |
| 22          | M1          | O.BL   |       | submesic/medium              | GL   | c3   | grassland with poplar                   |
| 23          | M1          | O.BL   |       | submesic/medium              | c3   |      | Canada buffaloberry/HWR Aw              |
| 24          | M2          | O.BL   |       | submesic/medium              | GL   |      | fescue grassland                        |
| 25          | F2          | O.HG   |       | subhydric/rich               | CS   |      | closed shrub                            |
| 26          | M2          | O.BL   |       | submesic/medium              | GL   |      | grazed fescue grassland                 |
| 27          | M2          | O.BL   |       | submesic/medium              | GL   |      | grazed fescue grassland                 |
| 28          | M2          | O.BL   |       | submesic/medium              | GL   |      | fescue grassland                        |
| 29          | F3          | O.HG   |       | subhydric/rich               | CS   | FG   | closed shrub and grassland              |
| 30          | M2          | O.BL   |       | submesic/medium              | GL   |      | fescue grassland                        |
| 31          | M1          | O.BL   | O.HG  | subhygric/rich               | f1   |      | balsam poplar Pb                        |
| 32          | F3          | O.HG   |       | subhydric/rich               | CS   |      | closed shrub                            |
| 33          | F3          | O.HG   |       | subhydric/rich               | CS   |      | closed shrub                            |
| 34          | C1          | O.BL   |       | submesic/medium              | GL   |      | grassland with poplar                   |
| 35          | C1          | O.BL   |       | submesic/medium              | GL   |      | grassland with poplar                   |
| 36          | C1          | O.BL   | O.HG  | mesic/medium                 | d2   |      | white meadowsweet PI                    |
| 37          | C1          | O.EB   |       | subxeric/poor                | a1   |      | limber pine/juniper Fd-Pf               |
| 38          | C1          | O.EB   |       | mesic/medium                 | d3   |      | white meadowsweet Sw                    |
| 39          | C1          | O.EB   |       | mesic/rich                   | e1   |      | thimbleberry/pine grass PI              |
| 40          | C1          | O.EB   |       | mesic/rich                   | e1   |      | thimbleberry/pine grass PI              |
| 41          | C1          | O.EB   |       | mesic/rich                   | e3   |      | thimbleberry/pine grass Sw              |
| 42          | C1          | O.EB   |       | subxeric/poor                | a1   |      | limber pine/juniper Fd-Pf               |
| 43          | C1          | O.EB   |       | subxeric/poor                | a1   |      | limber pine/juniper Fd-Pf               |
| 44          | C1          | O.EB   |       | mesic/rich                   | e1   |      | thimbleberry/pine grass PI              |
| 45          | C1          | O.EB   |       | submesic/medium              | c1   |      | Canada buffaloberry/HWR Fd              |
| 46          | M1          | O.BL   | O.HG  | mesic/medium                 | d1   |      | white meadowsweet Fd                    |
| 47          | M1          | O.BL   | O.HG  | submesic/medium              | GL   | c3   | grassland; with Canada buffaloberry/HWR |
| 48          | FM1         | O.BL   | O.HG  | mesic/rich                   | e2   |      | thimbleberry/ pine grass Aw             |
| 49          | M1          | O.BL   | O.HG  | submesic/medium              | GL   |      | grassland; with Canada buffaloberry/HWR |
| 50          | M1          | O.BL   |       | submesic/medium              | GL   |      | grassland; with Canada buffaloberry/HWR |
| 51          | M2          | O.BL   |       | submesic/medium              | GL   |      | fescue grassland                        |



| POL. | PM* | SOIL1 | SOIL2 | REGIMES         | VEG1 | VEG2 | VEGETATION DESCRIPTION **      |
|------|-----|-------|-------|-----------------|------|------|--------------------------------|
| 52   | F2  | O.HG  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 53   | M2  | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 54   | F3  | CA.BL |       | submesic/medium | GL   |      | grazed fescue grassland        |
| 55   | F4  | O.HG  | R.G   | subhydic/rich   | CS   | SM   | closed shrub; sedge meadows    |
| 56   | F4  | O.HG  | R.G   | subhydic/rich   | CS   |      | closed shrub                   |
| 58   | FM1 | O.HG  |       | subhydic/rich   | f1   |      | Balsam poplar Pb               |
| 59   | M2  | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 60   | FM1 | O.BL  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 61   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 62   | MC1 | O.BL  |       | subhydic/rich   | f1   |      | Balsam poplar Pb               |
| 63   | M2  | O.BL  |       | submesic/medium | GL   |      | grassland                      |
| 64   | MC1 | O.BL  |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw     |
| 65   | FM1 | O.HG  |       | subhydic/rich   | CS   | f1   | closed shrub; Balsam poplar Pb |
| 66   | FM1 | O.BL  | GL.BL | subhydic/rich   | f1   |      | Balsam poplar Pb               |
| 67   | FM1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 68   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 69   | MC1 | O.BL  |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw     |
| 70   | FM1 | O.HG  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 71   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 72   | MC1 | O.BL  |       | subhydic/rich   | f1   |      | Balsam poplar Pb               |
| 73   | FM1 | O.HG  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 74   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 75   | MC1 | O.BL  |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 76   | MC1 | O.BL  |       | mesic/medium    | d3   |      | white meadowsweet Sw           |
| 77   | F4  | O.HG  | R.G   | subhydic/rich   | CS   |      | closed shrub                   |
| 78   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 79   | MC1 | O.BL  |       | mesic/rich      | e1   |      | thimbleberry/ pine grass Pl    |
| 80   | MC1 | O.BL  |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 81   | FM1 | O.HG  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 82   | MC1 | O.BL  |       | mesic/rich      | e3   |      | thimbleberry/pine grass Sw     |
| 83   | FM1 | O.HG  |       | subhydic/rich   | CS   | SM   | grassland/shrubby              |
| 84   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 85   | MC1 | O.BL  |       | mesic/rich      | e2   |      | thimbleberry/pinegrass Aw      |
| 86   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 87   | MC1 | O.BL  |       | subxeric/medium | GL   |      | drier fescue grassland         |
| 88   | MC1 | O.BL  |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 89   | MC1 | O.BL  |       | mesic/rich      | e3   |      | thimbleberry/pine grass Sw     |
| 90   | MC1 | O.BL  | O.DG  | submesic/medium | GL   |      | fescue grassland               |
| 91   | MC1 | O.BL  | O.DG  | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 92   | MC1 | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 93   | MC1 | O.BL  | O.DG  | submesic/medium | GL   |      | fescue grassland               |
| 94   | MC1 | O.BL  | O.DG  | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 95   | MC1 | O.BL  |       | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw     |
| 96   | F1  | O.HG  | R.HG  | subhydic/rich   | f1   |      | Balsam poplar Pb               |
| 97   | F4  | O.HG  |       | subhydic/rich   | CS   |      | closed shrub                   |
| 98   | MC1 | O.EB  |       | submesic/medium | GL   |      | grazed grassland               |
| 99   | M2  | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 100  | MC1 | O.DG  | O.EB  | mesic/rich      | e2   |      | thimbleberry/pine grass Aw     |
| 101  | MC1 | O.DG  | O.EB  | mesic/rich      | e3   |      | thimbleberry/pine grass Sw     |
| 102  | M3  | O.EB  |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 103  | M2  | O.BL  |       | submesic/medium | GL   |      | fescue grassland               |
| 104  | FM1 | O.HG  |       | subhydic/rich   | CS   | f1   | closed shrub; balsam poplar Pb |
| 105  | MC1 | O.DG  | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland        |
| 106  | FM1 | O.HG  |       | subhydic/rich   | f1   | CS   | Balsam poplar Pb; closed shrub |
| 107  | MC1 | O.DG  | O.EB  | mesic/rich      | e3   |      | thimbleberry/pine grass Sw     |



| POL. | PM* | SOIL1  | SOIL2 | REGIMES         | VEG1 | VEG2 | VEGETATION DESCRIPTION **             |
|------|-----|--------|-------|-----------------|------|------|---------------------------------------|
| 108  | M3  | O.EB   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 109  | M3  | O.EB   |       | mesic/rich      | e1   |      | thimbleberry/pine grass PI            |
| 110  | FM1 | O.HG   | O.EB  | subhygric/rich  | f1   |      | balsam poplar Pb                      |
| 111  | M3  | O.EB   |       | mesic/rich      | e1   |      | thimbleberry/pine grass PI            |
| 112  | M3  | O.EB   |       | mesic/rich      | e1   |      | thimbleberry/pine grass PI            |
| 113  | F1  | O.HG   | R.HG  | subhygric/rich  | f1   |      | balsam poplar Pb                      |
| 114  | MC1 | O.BL   | O.DG  | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 115  | MC1 | O.BL   | O.DG  | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 116  | MC1 | O.BL   |       | submesic/medium | c3   | FG   | Canada buffaloberry/HWR Aw; grassland |
| 117  | C1  | O.EB   | O.DG  | mesic/rich      | e3   |      | thimbleberry/pine grass Sw            |
| 118  | MC1 | O.BL   | O.DG  | mesic/rich      | e1   |      | thimbleberry/pine grass PI            |
| 119  | MC1 | O.EB   | O.DG  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 120  | MC1 | O.BL   | O.DG  | mesic/rich      | e1   |      | thimbleberry/pine grass PI            |
| 121  | MC1 | O.BL   | O.DG  | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 122  | MC1 | O.BL   | O.DG  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 123  | MC1 | O.BL   | O.DG  | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 124  | M1  | O.BL   | O.DG  | mesic/rich      | e2   |      | thimbleberry/ pine grass Aw           |
| 125  | MC1 | O.BL   | O.DG  | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw            |
| 126  | MC1 | O.BL   |       | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw            |
| 127  | MC1 | O.BL   | O.DG  | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 128  | C2  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 129  | MC1 | O.BL   | O.DG  | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 130  | FM1 | O.HG   | O.BL  | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 131  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 132  | C2  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 133  | MC1 | O.BL   | O.DG  | mesic/rich      | e2   |      | thimbleberry/pinegrass Aw             |
| 134  | FM1 | O.BL   | O.DG  | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 135  | MC1 | O.BL   | O.DG  | submesic/medium | GL   |      | fescue grassland                      |
| 136  | MC1 | O.BL   | O.DG  | submesic/medium | GL   |      | fescue grassland                      |
| 137  | C2  | O.GL   | O.EB  | submesic/poor   | b2   |      | bearberry Aw                          |
| 138  | MC1 | O.BL   | O.DG  | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 139  | C2  | O.GL   | O.EB  | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 140  | MC1 | O.BL   | O.DG  | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 141  | CR1 | O.EB   | O.R   | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 142  | CR1 | O.EB   | O.R   | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 143  | CR1 | O.EB   | O.R   | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 144  | C1  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 145  | F1  | O.HG   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 146  | F1  | O.HG   |       | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 147  | MC1 | O.EB   |       | mesic/medium    | d1   |      | white meadowsweet Fd                  |
| 148  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 149  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 150  | F1  | O.HG   |       | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 151  | C2  | O.GL   | O.EB  | mesic/medium    | d1   |      | white meadowsweet Fd                  |
| 152  | C2  | O.GL   | O.EB  | submesic/poor   | b2   |      | bearberry Aw                          |
| 153  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf             |
| 154  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 155  | C2  | O.GL   | O.EB  | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 156  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf             |
| 157  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 158  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 159  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw                          |
| 160  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw                          |
| 161  | F1  | O.HG   |       | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw             |
| 162  | C2  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |



| POL. | PM* | SOIL1  | SOIL2 | REGIMES         | VEG1 | VEG2 | VEGETATION DESCRIPTION **  |
|------|-----|--------|-------|-----------------|------|------|----------------------------|
| 163  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 164  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 165  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 166  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 167  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 168  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 169  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 170  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 171  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 172  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 173  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 174  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 175  | C2  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI       |
| 176  | C1  | O.EB;  |       | mesic/medium    | d3   |      | white meadowsweet Sw       |
| 177  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI       |
| 178  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 179  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 180  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 181  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Aw |
| 182  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 183  | CR1 | O.EBsh |       | subxeric/poor   | GL   |      | grassland                  |
| 184  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 185  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI       |
| 186  | F1  | O.HG   | R.G   | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw  |
| 187  | C2  | O.GL   | O.EB  | submesic/medium | c2   |      | Canada buffaloberry/HWR PI |
| 188  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 189  | C2  | O.GL   | O.EB  | subxeric/medium | GL   |      | grassland                  |
| 190  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI       |
| 191  | F1  | O.HG   | R.G   | mesic/rich      | e3   |      | thimbleberry/pinegrass Sw  |
| 192  | C2  | O.GL   | O.EB  | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 193  | C2  | O.GL   | O.EB  | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 194  | C2  | O.GL   | O.EB  | mesic/medium    | d1   |      | white meadowsweet Fd       |
| 195  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI       |
| 196  | C2  | O.GL   | O.EB  | subhygric/rich  | f1   |      | balsam poplar Pb           |
| 197  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 198  | CR1 | O.EBsh |       | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw |
| 199  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 200  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 201  | C2  | O.GL   | O.EB  | submesic/poor   | b2   |      | bearberry Aw               |
| 202  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 203  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 204  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 205  | CR1 | O.EBsh |       | subxeric/poor   | GL   |      | grassland                  |
| 206  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 207  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 208  | C2  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland    |
| 209  | C2  | O.GL   | O.EB  | mesic/medium    | d2   |      | white meadowsweet PI       |
| 210  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 211  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 212  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |
| 213  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 214  | CR1 | O.EBsh |       | subxeric/poor   | GL   |      | dry grassland              |
| 215  | C2  | O.GL   | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd |
| 216  | CR1 | O.EBsh |       | subxeric/poor   | a1   |      | limber pine/juniper Fd-Pf  |
| 217  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw               |



| POL. | PM* | SOIL1  | SOIL2 | REGIMES         | VEG1 | VEG2 | VEGETATION DESCRIPTION **             |
|------|-----|--------|-------|-----------------|------|------|---------------------------------------|
| 218  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 219  | CR1 | O.EBsh |       | submesic/poor   | b2   |      | bearberry Aw                          |
| 220  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 221  | CR1 | O.EBsh |       | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw            |
| 222  | CR1 | O.EBsh |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 223  | C2  | O.GL   | O.EB  | submesic/medium | GL   |      | grassland                             |
| 224  | C2  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 225  | C2  | O.GL   | O.EB  | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 226  | M3  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 227  | M3  | O.EB   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 228  | M3  | O.EB   |       | mesic/rich      | e3   |      | thimbleberry/pine grass Sw            |
| 229  | M3  | O.EB   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 230  | M3  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 231  | M3  | O.EB   |       | submesic/medium | GL   |      | grassland                             |
| 232  | M3  | O.EB   |       | submesic/medium | c4   |      | Canada buffaloberry/HWR Aw-Sw-PI-Fd   |
| 233  | M3  | O.EB   |       | submesic/medium | c4   |      | Canada buffaloberry/HWR Aw-Sw-PI-Fd   |
| 234  | M3  | O.EB   |       | submesic/medium | GL   |      | grassland                             |
| 235  | M3  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 236  | M3  | O.EB   |       | submesic/medium | GL   |      | grassland                             |
| 237  | M3  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 238  | M1  | O.BL   |       | submesic/medium | GL   |      | grassland                             |
| 239  | M1  | O.BL   |       | submesic/medium | GL   |      | grassland                             |
| 240  | M1  | O.BL   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 241  | M1  | O.BL   |       | submesic/medium | c4   |      | Canada buffaloberry/HWR Aw-Sw-PI-Fd   |
| 242  | C1  | O.EB   |       | mesic/medium    | d3   |      | white meadowsweet Sw                  |
| 243  | C1  | O.EBsh |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 244  | C1  | O.EB   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 245  | C1  | O.EB   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 246  | C1  | O.EB   |       | mesic/rich      | e3   |      | thimbleberry/pine grass Sw            |
| 247  | C1  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 248  | MC1 | O.DG   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 249  | MC1 | O.DG   |       | submesic/medium | GL   |      | grassland                             |
| 250  | MC1 | O.DG   |       | mesic/rich      | e2   | FG   | thimbleberry/pine grass Aw; grassland |
| 251  | MC1 | O.DG   |       | submesic/medium | GL   |      | grassland                             |
| 252  | C1  | O.EB   |       | submesic/medium | c1   | c3   | Canada buffaloberry/HWR Fd, Aw        |
| 253  | C1  | O.EB   |       | submesic/medium | c3   | FG   | Canada buffaloberry/HWR Aw; grassland |
| 254  | C1  | O.EB   |       | submesic/medium | GL   |      | grassland                             |
| 255  | C1  | O.EB   |       | submesic/medium | c3   | c1   | Canada buffaloberry/HWR Aw, Fd        |
| 256  | F2  | O.HG   |       | subhydric/rich  | CS   |      | shrubland                             |
| 257  | MC1 | O.DG   | O.BL  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 258  | C1  | O.EB   |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 259  | C1  | O.EB   |       | submesic/medium | c3   |      | Canada buffaloberry/HWR Aw            |
| 260  | M1  | O.BL   | O.HG  | mesic/rich      | e2   |      | thimbleberry/pine grass Aw            |
| 261  | FM1 | O.HG   |       | mesic/rich      | e3   |      | thimbleberry/pine grass Sw            |
| 262  | M1  | O.BL   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 263  | C1  | O.EBsh |       | submesic/medium | GL   |      | fescue grassland                      |
| 264  | CR1 | O.R    | O.EB  | subxeric/poor   | a1   | FG   | limber pine/juniper Fd-Pf; grassland  |
| 265  | C1  | O.EBsh |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland               |
| 266  | C1  | O.EBsh |       | submesic/medium | c1   | c3   | Canada buffaloberry/HWR Fd; Aw        |
| 267  | M1  | O.BL   |       | mesic/medium    | d3   | d1   | white meadowsweet Sw (White spruce/mo |
| 269  | C1  | O.EB   |       | mesic/medium    | d3   | d1   | white meadowsweet Sw (White spruce/mo |
| 270  | C1  | O.EB   |       | mesic/medium    | d3   | d1   | white meadowsweet Sw (White spruce/mo |
| 271  | C1  | O.EB   |       | mesic/medium    | d2   |      | white meadowsweet PI                  |
| 272  | C1  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd            |
| 273  | C1  | O.EB   |       | submesic/poor   | GL   |      | grassland                             |



| POL. | PM* | SOIL1  | SOIL2 | REGIMES         | VEG1 | VEG2 | VEGETATION DESCRIPTION **            |
|------|-----|--------|-------|-----------------|------|------|--------------------------------------|
| 274  | C1  | O.EB   |       | submesic/poor   | b2   |      | bearberry Aw                         |
| 275  | C1  | O.EBsh |       | subxeric/poor   | a1   | FG   | limber pine/juniper Fd-Pf; grassland |
| 276  | C1  | O.EB   |       | subxeric/poor   | a1   | FG   | limber pine/juniper Fd-Pf; grassland |
| 277  | M1  | O.BL   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw           |
| 278  | M1  | O.BL   |       | submesic/medium | GL   |      | fescue grassland                     |
| 279  | C1  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd           |
| 280  | C1  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd           |
| 281  | CR1 | O.R    | O.EB  | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd           |
| 282  | M1  | O.BL   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw           |
| 283  | C1  | O.EB   |       | submesic/poor   | b2   | FG   | bearberry Aw; grassland              |
| 284  | M1  | O.BL   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw           |
| 285  | M1  | O.BL   |       | mesic/rich      | e2   |      | thimbleberry/pine grass Aw           |
| 286  | C1  | O.EB   |       | submesic/medium | c1   |      | Canada buffaloberry/HWR Fd           |

POL. = map polygon number

\* Refer to Appendix B for definitions of codes

\*\* refer to Archibald et al (1996) for further information on ecosite phases in forested polygons



APPENDIX B. Descriptions of physical land unit codes from Appendix A - Upper Bob Creek Ecological Reserve.

| PM<br>UNIT<br>CODE           | PARENT MATERIALS AND<br>LANDFORM                                                                                                                                                            | SOIL<br>CLASSIFICATION*                       | SLOPE<br>CLASS | DRAINAGE<br>CLASS             | VEGETATION - summary                                                                                                                                    |
|------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------|----------------|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>COLLUVIAL</b>             |                                                                                                                                                                                             |                                               |                |                               |                                                                                                                                                         |
| C1                           | Colluvial and residual veneers<br>over inclined or ridged bedrock;<br>upper slopes; strong to extreme<br>slopes with various aspects                                                        | O.EB - often<br>shallow to bedrock            | 6-7;<br>7-8    | Rapid                         | South-facing slopes are grassland or<br>open mixed forest with meadows;<br>north and west -facing are closed<br>forest of pine or douglas fir           |
| C2                           | Colluvial and residual veneers<br>over inclined eroded bedrock;<br>upper slopes                                                                                                             | O.GL; O.EB                                    | 6-7            | Well to .<br>rapid            | North-facing are closed forest (pine or<br>pine-spruce); south-facing are dry<br>grassland with aspen copses or<br>somewhat more open pine forest       |
| <b>COLLUVIAL AND BEDROCK</b> |                                                                                                                                                                                             |                                               |                |                               |                                                                                                                                                         |
| CR1                          | Colluvial veneers over inclined<br>eroded bedrock, with exposures<br>of ridged bedrock; shallow poorly<br>developed soils; occurs on or near<br>ridgetops; very strong to extreme<br>slopes | O.EB shallow to<br>bedrock (dominant);<br>O.R | 7-8;<br>8-9    | Rapid to<br>very rapid        | Open forest - limber pine-douglas fir or<br>aspen with meadows; or grassland<br>with scattered trees, some eroded<br>areas                              |
| <b>MORaine AND COLLUVIAL</b> |                                                                                                                                                                                             |                                               |                |                               |                                                                                                                                                         |
| MC1                          | Morainal and colluvial veneers<br>over bedrock; lower and mid<br>slopes west of Bob Creek valley                                                                                            | O.BL (dominant);<br>O.DG (subdominant)        | 5-6            | Well to<br>moderately<br>well | South-facing slopes are grassland with<br>some aspen woodland and some open<br>shrub; north-facing slopes are forested<br>(mixed or coniferous forests) |



| MORAINAL             |                                                                                                                                                                             |                                         |     |                           |                                                                       |
|----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|-----|---------------------------|-----------------------------------------------------------------------|
| M1                   | Rolling morainal veneers and blankets over inclined bedrock; lower slopes close to Bob Creek valley                                                                         | O.BL (dominant)<br>O.HG (minor amounts) | 5-6 | Moderately well           | Mixed forest with meadows; grasslands with scattered poplar woodlands |
| M2                   | Undulating till deposits on lower slopes close to bottom of Bob Creek valley                                                                                                | O.BL                                    | 3-4 | Moderately well           | Grasslands                                                            |
| M3                   | Morainal veneers over inclined eroded bedrock; mid-elevations                                                                                                               | O.EB                                    | 5-6 | Well to moderately well   | Coniferous forests of pine and douglas fir                            |
| FLUVIAL AND MORAINAL |                                                                                                                                                                             |                                         |     |                           |                                                                       |
| FM1                  | Morainal till deposits (dominant) interspersed with level fluvial deposits; lower slopes and ridges dissected by drainage channels;                                         | O.HG; O.BL; GL.BL                       | 2-3 | Imperfectly to poorly     | shrubby (willows; bog birch); also balsam poplar                      |
| FLUVIAL              |                                                                                                                                                                             |                                         |     |                           |                                                                       |
| F1                   | Fluvial veneers and blankets over inclined bedrock; Bottoms of coulees in upper slope areas;                                                                                | O.HG (dominant);<br>R.G (subdominant)   | 5   | Imperfectly to poorly     | Tall shrubs; or forested with spruce or spruce-pine                   |
| F2                   | Fluvial veneers over undulating morainal till deposits in fluvial terraces and other areas associated with older channels of Bob Creek, and in coulees in lower slope areas | O.HG (dominant);<br>CA.BL, CU.R         | 3   | Moderately well to poorly | Shrubby - most commonly willows                                       |
| F3                   | Bottom of Bob Creek valley; level fluvial deposits                                                                                                                          | O.HG (dominant);<br>R.G and CU.R        | 3   | Poorly to very poorly     | Grassy-sedge meadows and closed shrub (willows and bog birch)         |

\* codes for soil classification are from the Canadian System of Soil Classification (Canada Soil Survey 1987)



APPENDIX C. Summary of vegetation, landform and soils information collected at each site, Upper Bob Creek Ecological Reserve, 1994 and 1995. Refer to Fig. 1 for site locations.

a) grassland and shrubby lowland sites:

| Site | Vegetation community type*     | Parent materials**                         | Soil type**    |
|------|--------------------------------|--------------------------------------------|----------------|
| 01   | <i>Festcam-Festida-Dantpar</i> | Undulating moraine                         | O.BL           |
| 02   | <i>Festida-Dantpar-Festsca</i> | Morainal and residual veneers over bedrock | O.BL eroded    |
| 03   | <i>Festcam-Festida-Dantpar</i> | Morainal veneers                           | O.BL           |
| 04   | <i>Betugla-Salipet-Salimyr</i> | Morainal blankets                          | O.HG           |
| 05   | <i>Festcam-Festida-Dantpar</i> | Morainal veneers                           | O.BL           |
| 06   | <i>Festcam-Festida-Dantpar</i> | Morainal and residual veneers over bedrock | O.BL           |
| 07   | <i>Calarub-Elyminn</i>         | Colluvial veneers                          | O.R            |
| 08   | <i>Festcam-Festida-Dantpar</i> | Colluvial and residual veneers             | O.EBsh;<br>O.R |
| 09   | <i>Poaprat-Festsca</i>         | Fluvial veneers over undulating till       | CU.R;<br>CA.BL |
| 10   | <i>Festcam-Festida-Dantpar</i> | Morainal veneers                           | O.BL           |
| 11   | <i>Betugla-Salipet-Salimyr</i> | Fluvial deposits                           | O.HG           |
| 12   | <i>Festcam-Festida-Dantpar</i> | Morainal veneers                           | O.BL           |
| 13   | <i>Festcam-Festida-Dantpar</i> | Morainal veneers                           | O.BL           |
| 20   | <i>Festcam-Festida-Dantpar</i> | Colluvial veneers over bedrock             | O.EBsh         |
| 21   | <i>Betugla-Salipet-Salimyr</i> | Fluvial veneers over undulating moraine    | O.HG           |
| 22   | <i>Betugla-Salipet-Salimyr</i> | Fluvial veneers over undulating moraine    | O.HG           |
| 23   | <i>Festcam-Festida-Dantpar</i> | Morainal and colluvial veneers             | O.DG           |
| 24   | <i>Festcam-Festida-Dantpar</i> | Undulating moraine                         | O.BL           |



| Site | Vegetation community type*     | Parent materials**                          | Soil type** |
|------|--------------------------------|---------------------------------------------|-------------|
| 25   | <i>Poaprat-Festcam</i>         | Morainal and colluvial veneers              | O.DG/O.EB   |
| 26   | <i>Festsca-Festida-Dantpar</i> | Morainal and residual veneers               | O.BL        |
| 27   | <i>Festida-Dantpar-Festsca</i> | Morainal and colluvial veneers              | O.BL        |
| 28   | <i>Festcam-Festida-Dantpar</i> | Colluvial and residual veneers over bedrock | O.EBsh      |
| 29   | <i>Festsca-Poaprat-Phlepra</i> | Undulating moraine                          | O.BL        |

b) forested sites:

| Site | Vegetation community type*<br>(ecosite phase in brackets)**       | Surficial materials***                               | Soils***      |
|------|-------------------------------------------------------------------|------------------------------------------------------|---------------|
| 30   | <i>Picegla/moss</i><br>(white meadowsweet Sw)                     | Colluvial and residual veneers over inclined bedrock | O.EBsh        |
| 31   | <i>Poputre/Rosaaci/Calarub</i><br>(thimbleberry/pine grass Aw)    | Morainal and colluvial veneers over rock             | O.BL          |
| 32   | <i>Picegla/moss</i><br>(white meadowsweet Sw)                     | Colluvial and residual veneers over inclined bedrock | O.EBsh        |
| 33   | <i>Pseumen/Elyminn</i><br>(Canada buffaloberry/Hairy wild rye Fd) | Colluvial and residual veneers over inclined bedrock | O.EBsh        |
| 34   | <i>Poputre/Rosaaci/Calarub</i><br>(thimbleberry/pine grass Aw)    | Morainal veneers and blankets                        | O.BL;<br>O.HG |
| 35   | <i>Pseumen/Elyminn</i><br>(see #33)                               | Colluvial veneers over rock                          | O.GL;O.EB     |
| 36   | <i>Pseumen/Elyminn</i><br>(see #33)                               | Colluvial veneers over rock                          | O.GL;O.EB     |
| 37   | <i>Poputre-Picegla/Vaccsco</i>                                    | Till and colluvial veneers over rock                 | O.BL;O.DG     |
| 38   | <i>Poputre/Rosaaci/Calarub</i><br>(see #34)                       | Morainal blankets                                    | O.BL          |



| Site | Vegetation community type*<br>(ecosite phase in brackets)**    | Surficial materials***                                  | Soils***  |
|------|----------------------------------------------------------------|---------------------------------------------------------|-----------|
| 39   | <i>Pseumen/Elyminn</i><br>(see #33)                            | Colluvial and residual veneers<br>over inclined bedrock | O.EBsh    |
| 40   | <i>Pinucon/Spirbet</i><br>(white meadowsweet Pl)               | Morainal blankets                                       | O.EB      |
| 41   | <i>Picegla/moss</i><br>(see #32)                               | Morainal veneers and blankets                           | O.BL;O.HG |
| 42   | <i>Picegla-Pseumen/Spirbet</i><br>(white meadowsweet Fd)       | Colluvial and residual veneers<br>over inclined bedrock | O.EBsh    |
| 43   | <i>Picegla-Pseumen/Spirbet</i><br>(see #42)                    | Morainal and colluvial veneers                          | O.BL;O.DG |
| 44   | <i>Picegla/Rubupar</i><br>(thimbleberry/pine grass Sw)         | Morainal and colluvial veneers                          | O.BL      |
| 45   | <i>Picegla-Popubal/Sympalb</i><br>(balsam poplar Pb)           | Fluvial deposits                                        | O.HG      |
| 46   | <i>Poputre/Arctuva/Festsca</i><br>(bearberry Aw)               | Morainal and colluvial veneers                          | O.EB      |
| 47   | <i>Poputre/Rosaaci/Calarub</i><br>(thimbleberry/pine grass Aw) | Morainal blankets                                       | O.EB      |

\* seven-letter latin acronyms for plant species are from Alberta Environ. Protection (1993);

\*\* Ecosite phase descriptions (Archibald et al 1996): Aw - aspen, Fd - Douglas fir, Pf - limber pine, Pl - lodgepole pine, Pb - balsam poplar, Sw - white spruce

\*\*\* information on parent materials and soil types is from site data where available, or from Leskiw (1993)



APPENDIX D. Significant plant species which may occur in the Ecological Reserve, and status, habitat and occurrence details.

| Species of concern <sup>a</sup> :           |                             | Ranking<br>(ANHIC) |
|---------------------------------------------|-----------------------------|--------------------|
| <i>Erigeron radicans</i> <sup>b,c</sup>     | Dwarf fleabane              | S2                 |
| <i>Oryzopsis exigua</i> <sup>b,c</sup>      | Little rice grass           | S1                 |
| <i>Phacelia hastata</i> <sup>c</sup>        | Silver-leaved scorpion weed | S2                 |
| <i>Plantago canescens</i> <sup>c</sup>      | Western ribgrass            | S2                 |
| <i>Potentilla plattensis</i> <sup>b,c</sup> | Low cinquefoil              | S2                 |
| <i>Senecio cymbalarioides</i> <sup>c</sup>  | Alpine groundsel            | S1                 |
| <i>Spiraea densiflora</i> <sup>c</sup>      | Pink meadowsweet            | S2                 |
| <b>Other significant species</b>            |                             |                    |
| <i>Goodyera oblongifolia</i> <sup>c</sup>   | Rattlesnake plantain        | --                 |
| <i>Microseris nutans</i> <sup>c</sup>       | Nodding scorzonella         | S2S3               |

<sup>a</sup> Alberta Natural Heritage Information Centre: Tracking list of plant species of special concern.

<sup>b</sup> Species of concern which has been found nearby (on Black Mountain east of the study area)

<sup>c</sup> Recorded by Brown et al (1986)

1) *Erigeron radicans* Hook. (Dwarf fleabane)

**Status:** species of special concern ranked S2 on tracking list (ANHIC 1996); rare in Alberta (Packer and Bradley 1984, Wallis et al 1987); rare in Canada (Fairbairns et al 1987); western species, rare throughout its range (Wallis et al 1986)

**Habitat:** dry ridges, open slopes, scree slopes (mostly north of Upper Bob Creek Ecological Reserve)

**Occurrence:** recorded by Wallis et al (1986) at the south end of Whaleback Ridge, and on west-facing slope of nearby Black Mountain

2) *Goodyera oblongifolia* Raf. (Rattlesnake plantain)

**Status:** uncommon in Alberta; scattered records in the foothills (Moss 1983)

**Habitat:** woods

**Occurrence:** recorded by Brown et al (1986)



3) *Microseris nutans* (Hook.) Sch.-Bip. (Nodding scorzonella)

**Status:** rare in Alberta (Packer and Bradley 1984; Wallis et al 1987); southwest Alberta distribution

**Habitat:** dry to moist slopes, open woods; Douglas fir stands (Wallis et al 1987);

**Occurrence:** recorded by Brown et al (1986)

4) *Oryzopsis exigua* Thurb. (Little rice grass)

**Status:** rare in Alberta (Packer and Bradley; ; Wallis et al 1987); rare in Canada (Fairbarns et al 1987); distribution is southwest corner of the province

**Habitat:** dry open ground or open woods

**Occurrence:** recorded by Brown et al (1986)

5) *Phacelia hastata* Doug. ex Lehm. (Silver-leaved scorpion weed)

**Status:** uncommon in Alberta; in southwest part of province (Moss 1983); species of concern with only 6-20 occurrences (ANHIC 1996)

**Habitat:** dry slopes and valleys

**Occurrence:** recorded by Brown et al (1986)

6) *Plantago canescens* Adams (Western ribgrass)

**Status:** uncommon in Alberta; 7 records in the province (Moss 1983); species of concern with only 6-20 occurrences (ANHIC 1996)

**Habitat:** grassy and gravelly slopes

**Occurrence:** recorded by Brown et al (1986)

7) *Potentilla plattensis* Nutt. (Low cinquefoil)

**Status:** rare in Alberta (Packer and Bradley 1984); only 3 records; species of fescue grasslands

**Habitat:** dry areas in grassland and coulees

**Occurrence:** recorded by Brown et al (1986)

8) *Senecio cymbalarioides* Buck (Alpine groundsel)

**Status:** rare in Alberta (Packer and Bradley 1984; Wallis et al 1987); rare in Canada (Fairbarns et al 1987); southwestern Alberta species

**Habitat:** moist subalpine meadows; 2000-2300 m;

**Occurrence:** recorded by Brown et al (1986)

9) *Spiraea densiflora* Nutt. (Pink meadowsweet)

**Status:** rare in Alberta (Packer and Bradley 1984)


**Habitat:** moist meadows, thickets and woods, boggy places (4)

**Occurrence:** recorded by Brown et al (1986)



APPENDIX E. Ranking system used to evaluate status of significant plant species. The system was originally developed by The Nature Conservancy, and is presented in ANHIC (1996). Only the ranks for species discussed in this report are included here.


| Provincial Rank | Definition                                                                                                                                                                                              |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| S1              | Five or fewer occurrences or only a few remaining individuals or may be imperiled because some factor of its biology makes it especially vulnerable to extirpation.                                     |
| S2              | 6-20 occurrences or with many individuals in fewer occurrences; or may be susceptible to extirpation because of some factor of its biology.                                                             |
| S3              | 21-100 occurrences, may be rare and local throughout its range, or in a restricted range (may be abundant in some locations or may be vulnerable to extirpation because of some factor of its biology). |
| SU              | Status uncertain often because of low search effort or cryptic nature of the element; possibly in peril                                                                                                 |



# UPPER B ECOLOGIC SIGNIFICANT VEG

WEST OF TH

SCALE



Prepared by  
Corporate Management Service;

Date of

## PLANIMETRIC SYMBOLS

- .....Hard Surface - All Weather
- .....Loose or Stabilized Surface - All Weather
- .....Loose Surface - Dry Weather
- .....Truck Trail
- .....Seismic Line or Trail
- .....Bridge
- .....Road Allowance
- .....Railway
- .....Transmission Line

DITCH

x-H

x

x





|       |                             |       |       |                                 |
|-------|-----------------------------|-------|-------|---------------------------------|
| ..... | Loose Surface - Dry Weather | DITCH | ..... | Ditch, Canal or Flume           |
| ..... | Truck Trail                 | ..... | ..... | Perennial Lake - Open Water     |
| ..... | Seismic Line or Trail       | ..... | ..... | Intermittent Lake               |
| ..... | Bridge                      | ..... | ..... | Gravel Bar, Sand or Mud         |
| ..... | Road Allowance              | ..... | ..... | Gate                            |
| ..... | Railway                     | ..... | ..... | Fence existing                  |
| ..... | Transmission Line           | ..... | ..... | Fence proposed                  |
| ..... | Pipeline                    | ..... | ..... | Map Unit Boundaries             |
| ..... | Oil Well site               | X 29  | ..... | Sample Sites                    |
| ..... | Gas Well site               | ..... | ..... | Grazing Lease / Permit Boundary |


# **SIGNIFICANT VEGETATION FEATURES**

|        |                        |        |                                      |
|--------|------------------------|--------|--------------------------------------|
| A..... | Angelica dawsonii      | E..... | Corallorhiza striata                 |
| B..... | Carex scopulorum       | F..... | Osmorhiza occidentalis               |
| C..... | Castilleja cusickii    | G..... | Poa nevadensis                       |
| D..... | Conimitella williamsii | H..... | Senecio foetidus var. hydrophiloides |

# UPPER BOB CREEK ECOLOGICAL RESERVE ECOLOGICAL LAND CLASSIFICATION

WEST OF THE FIFTH MERIDIAN

SCALE 1:15 000



Prepared by Resource Data Division  
Corporate Management Service; Alberta Environmental Protection; January 1997


Date of Orthophoto 1982

## VEGETATION CODES

| CODE + | DESCRIPTION ++                                 |
|--------|------------------------------------------------|
| a1     | Limber pine/juniper Fd                         |
| b2     | Bearberry Aw                                   |
| c1     | Canada buffaloberry/hairy wild rye Fd          |
| c2     | Canada buffaloberry/hairy wild rye Pl          |
| c3     | Canada buffaloberry/hairy wild rye Aw          |
| c4     | Canada buffaloberry/hairy wild rye Aw-Sw-Pl-Fd |
| d1     | White meadowsweet Fd                           |
| d2     | White meadowsweet Pl                           |
| d3     | White meadowsweet Sw                           |
| e1     | Thimbleberry/pinegrass Pl                      |
| e2     | Thimbleberry/pinegrass Aw                      |
| e3     | Thimbleberry/pinegrass Sw                      |
| fi     | Balsam poplar Pb                               |
| gl     | Grassland                                      |
| sm     | Sedge Meadow                                   |
| cs     | Closed shrub                                   |

### LABEL CODES


Polygon ID Numbers  
203  
a1-CR1  
Physical Land Unit Codes  
Vegetation Codes



+ Refer to Archibald et al (1996) for details of forest ecosite phases

++ Key to overstory species abbreviation

Aw - Aspen



sm Seede Meadow  
cs Closed shrub

+ Refer to Archibald et al(1996) for details of forest ecosite phases

++ Key to overstory species abbreviation

Aw - Aspen  
Fd - Douglas fir  
Pb - Balsam poplar  
Pf - Limber pine  
Pl - Lodgepole pine  
Sw - White spruce

## PHYSICAL LAND UNIT CODES

| PM UNIT CODE          | PARENT MATERIALS AND LANDFORM                                                                                                                                                | SOIL CLASSIFICATION*                        | SLOPE CLASS | DRAINAGE CLASS            | VEGETATION - summary                                                                                                                           |
|-----------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|-------------|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| COLLUVIAL             |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| C1                    | Colluvial and residual veneers over inclined or ridged bedrock; upper slopes; strong to extreme slopes with various aspects.                                                 | O.EB - often shallow to bedrock             | 6-7;<br>7-8 | Rapid                     | South-facing slopes are grassland or open mixed forest with meadows; north and west-facing are closed forest of pine and douglas fir           |
| C2                    | Colluvial and residual veneers over inclined eroded bedrock; upper slopes                                                                                                    | O.GL; O.EB                                  | 6-7         | Well to Rapid             | North-facing are closed forest (pine or pine-spruce); south-facing are dry grassland with aspen copees or somewhat more open pine forest       |
| COLLUVIAL AND BEDROCK |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| CR1                   | Colluvial veneers over inclined eroded bedrock; with exposures of ridged bedrock; shallow poorly developed soils; occurs on or near ridgetops; very strong to extreme slopes | O.EB shallow to bedrock (dominant); O.R     | 7-8;<br>8-9 | Rapid to very rapid       | Open forest-limber pine-douglas fir or aspen with meadows; or grassland with scattered trees, some eroded areas                                |
| MORaine AND COLLUVIAL |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| MC1                   | Morainel and colluvial veneers over bedrock; lower and mid slopes west of Bob Creek valley                                                                                   | O.BL (dominant); O.DG (subdominant)         | 5-6         | Well to moderately well   | South-facing slopes are grassland with some aspen woodland and some open shrub; north-facing slopes are forested (mixed or coniferous forests) |
| MORAINAL              |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| M1                    | Rolling morainel veneers and blankets over inclined bedrock; lower slopes close to Bob Creek valley                                                                          | O.BL (dominant)<br>O.HG (minor amounts)     | 5-6         | Moderately well           | Mixed forest with meadows; grasslands with scattered poplar woodlands                                                                          |
| M2                    | Undulating till deposits on lower slopes close to bottom of Bob Creek valley                                                                                                 | O.BL                                        | 3-4         | Moderately well           | Grassland                                                                                                                                      |
| M3                    | Morainel veneers over inclined eroded bedrock; mid-elevations                                                                                                                | O.EB                                        | 5-6         | Well to moderately well   | Coniferous forest of pine and douglas fir                                                                                                      |
| FLUVIAL AND MORAINAL  |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| FM1                   | Morainel till deposits (dominant) interspersed with level fluvial deposits; lower slopes and ridges dissected by drainage channels                                           | O.HG; O.BL; GL.BL                           | 2-3         | Imperfectly to poorly     | Shrubby (willow; bog birch); also balsam poplar                                                                                                |
| FLUVIAL               |                                                                                                                                                                              |                                             |             |                           |                                                                                                                                                |
| F1                    | Fluvial veneers and blankets over inclined bedrock; Bottoms of coulees in upper slopes areas                                                                                 | O.HG (dominant); R.G (subdominant)          | 5           | Imperfectly to poorly     | Tall shrubs, or mixed forest with white spruce or balsam poplar.                                                                               |
| F2                    | Fluvial veneers over undulating morainel till deposits in fluvial terraces and other areas associated with older channels of Bob Creek, and in coulees in lower slope areas  | O.HG (dominant); CABL, CU.R                 | 3           | Moderately well to poorly | Shrubby - most commonly willows                                                                                                                |
| F3                    | Bottom of Bob Creek valley; level fluvial deposits                                                                                                                           | O.HG (dominant); R.G and CU.R (subdominant) | 3           | Poorly to very poorly     | Grassland-sedge meadows and closed shrub (willows and bog birch)                                                                               |

\* codes for soil classification are from the Canadian System of Soil Classification (Canada Soil Survey 1987)







VE  
ON

METRE

MILE



Canada buffaloberryhaity wild rye P  
Canada buffaloberryhaity wild rye A

c3

c2



National Library of Canada  
Bibliothèque nationale du Canada



3 3286 51349 2146